

Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 0 819 758 A2**

(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:
21.01.1998 Bulletin 1998/04

(51) Int. Cl.⁶: **C12N 15/12**, C12N 15/24,
A61K 48/00

(21) Application number: **97112154.6**

(22) Date of filing: **16.07.1997**

(84) Designated Contracting States:
**AT BE CH DE DK ES FI FR GB GR IE IT LI LU MC
NL PT SE**
Designated Extension States:
AL LT LV RO SI

(30) Priority: **16.07.1996 US 680845**

(71) Applicant:
**Mixson, Archibald James
Rockville, MD 21201 (US)**

(72) Inventor:
**Mixson, Archibald James
Rockville, MD 21201 (US)**

(74) Representative:
**Sleckmann, Ralf, Dr. et al
Cohausz Hase Dawidowicz & Partner,
Patent- und Rechtsanwaltskanzlei,
Schumannstrasse 97
40237 Düsseldorf (DE)**


Remarks:

The applicant has subsequently filed a sequence listing and declared, that it includes no new matter.

(54) **Cationic vehicle: DNA complexes and their use in gene therapy**

(57) Cationic vehicles: DNA complexes comprising DNA encoding an anti-angiogenic peptide or DNA encoding a tumor suppressor protein and DNA encod-

ing an anti-angiogenic peptide, as well as their use in gene therapy, are disclosed.



*- BAP vs BAP-p53 or BAP-TSPf, $p < 0.05$

**BAP-p53 or BAP-TSPf vs BAP-p53/BAP-TSPf, $p < 0.01$

EP 0 819 758 A2

DescriptionFIELD OF THE INVENTION

5 The present invention relates to cationic vehicles DNA complexes (*i.e.* cationic liposome:DNA complexes, cationic polymer:DNA complexes) comprising DNA encoding an anti-angiogenic peptide, or DNA encoding a tumor suppressor protein and DNA encoding an anti-angiogenic peptide, and their use in gene therapy.

BACKGROUND OF THE INVENTION

10 I. Gene Therapy

Development of gene therapy techniques is approaching clinical realization for the treatment of neoplastic and metabolic diseases. The main obstacle in the treatment of malignant diseases, however, remains in the vector delivery system of the transgene to a distant target tissue.

15 Vectors carrying genes are commonly divided into viral and non-viral vector categories. Unfortunately, all vectors described to date have significant limitations. For example, replication-deficient retroviral vectors can efficiently transfect dividing cells. Local intratumoral injection of retroviruses that contain a thymidine kinase transgene has been used successfully to affect regression of gliomas (Culver et al, *Science*, 256:1550-1552 (1992)). However, retroviruses have the potential to cause insertional mutagenesis. As a result, their use has been limited to either direct injection of tumors or to *ex vivo* gene transfer trials. Unlike retroviral vectors, adenoviral vectors can also transfect non-dividing cells, and their ability to cause insertional mutagenesis is greatly reduced. However, they have the undesirable potential to activate the immune system in humans (Crystal, *Science*, 270:404-410, (1995). Attempts are underway to minimize the immunogenicity of the adenoviral vectors, but the potential toxicity of viral vectors will most likely limit their use for systemic delivery of genes in the near future.

25 Non-viral vectors of DNA include liposomes, peptides, proteins and polymers (Ledley, *Current Opinion in Biotechnology*, 5:626-636 (1994)). Of these, liposomes are the most commonly used non-viral vectors of DNA. The major advantage of liposomes over retroviruses is that DNA is not incorporated into the genome, and unlike adenoviral vectors, they are not immunogenic. However, the major limitation of liposomes is that they are not as efficient as viral vectors in transfecting many cell types. Until recently, their medical utility was limited by their rapid uptake by phagocytic cells. Interest in liposomes as a vector was rejuvenated by two technological advances that have produced a renaissance in the field. First, stearily stabilized (Stealth) liposomes represent a significant breakthrough in that they are non-reactive, and are not readily taken up by the reticuloendothelial system (RES). Stealth liposomes are composed of lipids rich in oxygen in their head group (ethylene glycol or glycolipids) which provide a steric barrier outside of the membrane. As a result, Stealth liposomes remain in the blood for up to 100 times longer than conventional liposomes, and can thus increase pharmacological efficacy (Papahadjopoulos, *In: Stealth liposomes*, Ed., Lasic et al, CRC Press (1995); and Lasic et al, *Science*, 267:1275-76 (1995)). However, Stealth liposomes are still not particularly efficient in transfection of cells or as vectors for DNA.

40 The second significant advance in liposome technology has been the use of cationic liposomes complexed to negatively-charged DNA. Cationic liposomes can condense DNA, and increase transfection yields several orders of magnitude. In the cationic liposome:DNA complex, the nucleic acids or oligonucleotides are not encapsulated, but are simply complexed with small unilamellar vesicles by electrostatic interactions. The exact nature of the cationic liposome:DNA complex is not clear, but intricate topological rearrangements of the cationic liposome:DNA complex occur, including DNA condensation, liposome aggregation, and fusion. This supramolecular complex can be added to cells *in vitro*, injected parenterally, or aerosolized for pulmonary applications (Lasic et al, *Science*, 267:1275-1276 (1995)). Further, the intravenous injection into mice of high concentrations of the CAT gene (100 µg or greater) complexed with cationic liposomes has been found to result in 40% transfection efficiency of well vascularized tissues, such as the spleen (Zhu et al, *Science*, 261:209-211 (1993)). However, a major challenge of gene therapy remains the systemic delivery of transgenes to the tumor or peritumoral area that will effectively decrease the size of primary tumors and their metastases. This is because unlike the spleen and bone marrow, which are highly vascular and have a high capacity to filter macromolecules from the blood stream, most organs and tumors do not have this capacity, and the transfection efficiency of these tissues with liposomes is low (Marshall, *Science*, 269:1051-1055 (1995)). In addition, another limitation of cationic liposome: DNA complexes is that their 1/2 life in the blood stream is less than one hour (Allen et al, *In: Liposome Technology-Vol. III*, Ed., Gregoriadis G et al, CRC Press (1993)). Sufficient transfection of the target cell by vectors carrying therapeutic genes has thus far been the rate-limiting step in gene therapy.

II. Tumor Suppressor Genes

Tumor suppressor genes are well-known in the art, and include the p53 gene (Baker et al, *Science*, 249:912-915 (1990)), the p21 gene (El-Deiry et al, *Cell*, 75:817-825 (1993); and Harper et al, *Cell*, 75:805-816 (1993)), and the rb gene (Bookstein et al, *Science*, 247:712-715 (1990)).

Mutations in the tumor suppressor gene p53 are known to occur in over 50% of human tumors, including metastatic breast cancer (Vogelstein, *Nature*, 348:681-682 (1990)). Breast cancer is one of the leading causes of death in women in North America and Western Europe, affecting nearly 10% of this population living to 80 years of age, and one million new cases are predicted by the end of this decade (Miller et al, *Int. J. Cancer*, 37:173-177 (1986)). Although the molecular basis of multistage carcinogenesis in breast cancer is not well understood, the metastatic potential of breast cancers has been correlated with the presence of point mutations in the p53 gene (Wang et al, *Oncogene*, 8:279-288 (1993)). Various groups have found that reintroduction of the wild-type P53 into a tumor cell has the therapeutic potential to inactivate the proliferative effects of the mutated product (Bookstein et al, *Cancer*, 71:1179-1186 (1993); Chen et al, *Science*, 250:1576-1580 (1990); and Baker et al, *Science*, 249:912-915 (1990)). For example, *in vitro* transfection and retroviral-mediated transfer of a single copy of the p53 transgene into a variety of tumor cells, including breast cancer cells, was found to result in a decrease in growth rate and/or attenuated tumor development once those transfected cells were implanted into nude mice (Wang et al, *Oncogene*, 8:279-288 (1993); Baker et al, *Science*, 249:912-915 (1990); Bookstein et al, *Science*, 247:712-715 (1990); Cheng et al, *Cancer Res.*, 52:222-226 (1992); Isaacs et al, *Cancer Res.*, 51:4716-4720 (1991); Diller et al, *Mol. Cell. Biol.*, 10:5772-5781 (1990); Chen et al, *Oncogene*, 6:1799-1805 (1991); and Zou et al, *Science*, 263:526-529 (1994)). In addition, intratracheal injection of a retrovirus containing the p53 transgene has been shown to significantly inhibit the growth of lung tumors (Fujiwara et al, *J. Natl. Cancer Inst.*, 86:1458-1462 (1994)). Further, systemic intravenous administration of a β -actin promoter-containing vector containing the p53 coding sequence complexed to cationic liposomes has been found to affect the tumor growth of a malignant line of breast cancer cells injected into nude mice (Lesoon-Wood et al, *Proc. Am. Ass. Cancer Res.*, 36:421 (1995); and Lesoon-Wood et al, *Human Gene Ther.*, 6:39-406 (1995)). Of the 15 tumors treated in this study, four of these tumors did not respond to treatment. Because of the unresponsiveness of these tumors, new therapies were sought in the present invention to more effectively decrease the size of these tumors. Based on the *in vitro* data concerning p53, one might expect that p53 decreases the size of the tumors due to efficient transfection of the tumor. However, less than 5% of the tumor was transfected after three injections of a cationic liposome:marker (CAT) gene. Furthermore, some endothelial cells of the tumor were transfected with this marker gene. Thus, the primary target of cationic liposome:p53 complex may be the vasculature system of the tumor. Given that angiogenesis is critical for the development of any human tumor, as well as for metastases (Fidler et al, *Cell*, 79:185-188 (1994)), this therapy should be widely applicable to a wide variety of tumors.

p53 coordinates multiple responses to DNA damage. DNA damage results in an increase in the level of the p53 protein. Following DNA damage, an important function of wild-type p53 function is to control the progression of cells from G1 to S phase. Recently, several groups have found that p53 transcriptionally activates a p21 kd protein (also known as WAF1 or Cip1), an inhibitor of cyclin-dependent kinases (CDKs) (El-Deiry et al, *supra*; and Harper et al, *supra*). Inhibition of CDK activity is thought to block the release of the transcription factor E2F, and related transcription factors from the retinoblastoma protein RB, with consequent failure to activate transcription of genes required for S phase entry (Harper et al, *supra*; and Xiong et al, *Nature*, 366:701-704 (1993)). Evidence consistent with the model that pRb is a downstream effector of p53-induced G1 arrest has recently been reported (Dulic et al, *Cell*, 76:1013-1023 (1994)). Thus, p53 regulates cell cycle through two proteins: p21 and rb.

III. Anti-Angiogenic Proteins

Proteins with anti-angiogenic activities are well-known and include: thrombospondin I (Kosfeld et al, *J. Biol. Chem.*, 267:16230-16236 (1993); Tolsma et al, *J. Cell Biol.*, 122:497-511 (1993); and Dameron et al, *Science*, 265:1582-1584 (1995)), IL-12 (Voest et al, *J. Natl. Cancer Inst.*, 87:581-586 (1995)), protamine (Ingber et al, *Nature*, 348:555-557 (1990)), angiostatin (O'Reilly et al, *Cell*, 79:315-328 (1994)), laminin (Sakamoto et al, *Cancer Res.*, 5:903-906 (1991)), and a prolactin fragment (Clapp et al, *Endocrinol.*, 133:1292-1299 (1993)). In addition, several anti-angiogenic peptides have been isolated from these proteins (Maione et al, *Science*, 247:77-79 (1990); Woltering et al, *J. Surg. Res.*, 50:245-251 (1991); and Eijan et al, *Mol. Biother.*, 3:38-40 (1991)).

Thrombospondin I (hereinafter "TSPI") is a large trimeric glycoprotein composed of three identical 180 kd subunits (Lahav et al, *Semin. Thromb. Hemostasis*, 13:352-360 (1987)) linked by disulfide bonds (Lawer et al, *J. Cell Biol.*, 103:1635-1648 (1986); and Lahav et al, *Eur. J. Biochem.*, 145:151-156 (1984)). The majority of anti-angiogenic activity is found in the central stalk region of this protein (Tolsma et al, *supra*). There are at least two different structural domains within this central stalk region that inhibit neovascularization (Tolsma et al, *supra*).

Besides TSPI, there are five other proteins (fibronectin, laminin, platelet factor-4, angiostatin, and prolactin frag-

ment) in which peptides have been isolated that inhibit angiogenesis. In addition, analogues of the peptide somatostatin are known to inhibit angiogenesis.

Fibronectin (FN) is a major surface component of many normal cells, as well as a potent cell spreading factor. During transformation, the loss of cellular FN has been observed. Furthermore, the addition of fibronectin to transformed cells restores the normal phenotype. It has been found that either heparin-binding or cell-adhesion fragments from FN can inhibit experimental metastasis, suggesting that cell surface proteoglycans are important in mediating the adhesion of metastatic tumor cells (McCarthy et al, *J. Natl. Cancer Inst.*, **80**:108-116 (1988)). It has also been found that FN and one of its peptides inhibits *in vivo* angiogenesis (Eijan et al, *Mol. Biother.*, **3**:38-40 (1991)).

Laminin is a major component of the basement membrane, and is known to have several biologically active sites that bind to endothelial and tumor cells. Laminin is a cruciform molecule that is composed of three chains, an A Chain and two B chains. Several sites in laminin have been identified as cell binding domains. These sites promote cellular activities *in vitro*, such as cell spreading, migration, and cell differentiation. Two peptides from two sites of the laminin B1 chain are known to inhibit angiogenesis (Grant et al, *Path. Res. Pract.*, **190**:854-863 (1994)).

Platelet factor-4 (PF4) is a platelet α -granule protein originally characterized by its high affinity for heparin. The protein is released from platelets during aggregation as a high molecular weight complex of a tetramer of the PF4 polypeptide and chondroitin sulfate, which dissociates at high ionic strength. PF4 has several biological properties including immunosuppression, chemotactic activity for neutrophils and monocytes as well as for fibroblasts, inhibition of bone resorption, and inhibition of angiogenesis. The angiostatic properties of human PF4 are associated with the carboxyl-terminal, heparin binding region of the molecule. A 12 amino acid synthetic peptide derived therefrom has been discovered to have marked angiostatic effects (Maione et al, *Science*, **247**:77-79 (1990)).

Although somatostatin is not a protein, it is a naturally-occurring cyclic 14 amino acid peptide whose most-recognized function is the inhibition of growth hormone (GH) secretion. Somatostatin is widely distributed in the brain, in which it fulfills a neuromodulatory role, and in several organs of the gastrointestinal tract, where it can act as a paracrine factor or as a true circulating factor. The role played by the neuropeptide somatostatin, also known as somatotropin release inhibitory factor (SRIF), in human cancer is not well understood. Recent investigations involving somatostatin receptors in normal and neoplastic human tissues suggest that the action is complex, and involves both direct and indirect mechanisms. One of the anti-tumor mechanisms of these synthetic somatostatin analogues may be an anti-angiogenic effect (Woltering et al, *J. Surg. Res.*, **50**:245-50 (1990)). In a recent study, the ability of native somatostatin and nine somatostatin analogues to inhibit angiogenesis were evaluated. The most potent somatostatin analogue was found to be approximately twice as potent as the naturally-occurring somatostatin (Barrie et al, *J. Surg. Res.*, **55**:446-50 (1993)).

Angiostatin is a 38 kDa polypeptide fragment of plasminogen. Whereas plasminogen has no fibrinogenic activity, angiostatin has marked angiogenic activity (O'Rielly MS, et al *Cell*, **79**:315-28 (1994)). Angiostatin was isolated when it was observed that the primary tumor suppressed metastases. That is, when the primary tumor was removed, the metastases grew. Administration of angiostatin blocks neo-vascularization and growth of metastases.

Finally, a 16kd fragment of prolactin has been found to be angiogenic. Similar to plasminogen, prolactin is not anti-angiogenic but the prolactin fragment is a potent *in vivo* and *in vitro* inhibitor of angiogenesis (Clapp C. et al. *Endocrinology*. **133**:1292-1299 (1993)).

Despite the evidence that anti-angiogenic peptides are effective anti-tumor agents, as well as the great deal of interest in targeting genes toward the vasculature, there have been no published reports on effective *in vivo* gene therapy regimens with established anti-angiogenic DNA sequences.

There are several reasons why gene therapy utilizing antiangiogenic genes have not been used or why antiangiogenic peptides are effective and the liposome: antiangiogenic gene may not be. First, there are significant physical differences between the liposome: DNA complexes and their peptides. Cationic liposomes have a 1/2 life of less than one hour (Allen TM and Papahajopoulos D, In: *Liposome Technology-Vol. III*, Ed., Gregoriadis G et al, CRC Press (1993)). whereas the most effective of the antiangiogenic peptides (i.e angiostatin) have a 1/2 life of two days (Folkman J, The John Krantz, Jr Lecture in Pharmacology, UMAB, 4/30/96). Since cationic liposomes form large aggregates when mixed with DNA, the distribution of these complexes is likely to be quite different from the much smaller peptides (need reference). These properties of the liposomes may account for the low transfection efficiency of a tumor. Therefore, it is uncertain as to whether these liposome:DNA complexes will reach their cellular targets.

Furthermore, the exact receptor target or mechanisms of these antiangiogenic peptides are unknown (Tolsma et al, *supra*). For example, it is unknown whether these receptor targets are intracellular or extracellular. The anti-angiogenic genes that are complexed to liposomes encode their respective proteins inside the cell, and proteins without secretory sequences remain inside the cell. Thus, it is unclear that a intracellular antiangiogenic peptide derived from a systemically transfected gene will reach its cellular and/or receptor target.

The only transfected antiangiogenic gene that has inhibited tumor growth is full length thrombospondin I. In this study (Weinstat-Saslow et al, *Cancer Research* **54**, 6504-6511, (1994)) tumor cells that expressed 15 fold higher levels of the thrombospondin I *in vitro* than baseline cells were implanted into the mice. This transfected full length throm-

bospondin I was secreted from the tumor cells to inhibit angiogenesis, and effectively reduced the tumor by 60%. Thus, this study determined that transfection of 100% of the tumor cells with a highly expressed secreted antiangiogenic gene was able to reduce tumor size.

5 SUMMARY OF THE INVENTION

An object of the present invention is to provide cationic vehicles:DNA complexes, such as liposome complexes containing DNA encoding anti-angiogenic peptides or cationic complexes containing DNA encoding an anti-angiogenic peptides.

10 Another object of the present invention is to provide a method of anti-angiogenic gene therapy.

Still another object of the present invention is to provide liposome complexes containing DNA encoding an anti-angiogenic peptide.

Yet another object of the present invention is to provide liposome complexes containing DNA encoding an anti-angiogenic peptide in combination with DNA encoding a tumor suppressor gene.

15 A further object of the present invention is to provide liposome complexes containing DNA encoding concatamers of the same or different anti-angiogenic peptides.

An additional object of the present invention is to provide a method for inhibiting tumor growth in a subject, or alternatively, to use these complexes for the production of a medicament, especially for inhibiting tumor growth in a subject.

20 These and other objects of the present invention, which will be apparent from the detailed description of the invention provided hereinafter, have been met in one embodiment, by a cationic liposome:DNA complex comprising DNA encoding an anti-angiogenic peptide and DNA encoding a tumor suppressor protein.

Further objects of the present invention were solved by the features defined in the present set of claims, but especially the preferred embodiments defined in claims 2 to 25.

25 Based on the present invention, it is anticipated that one skilled in the gene therapy could utilize other cationic carriers (polylysine, polyhistidine, polycat57, Superfect, and polyethylimine) complexed with the antiangiogenic genes to inhibit tumors.

DETAILED DESCRIPTION OF THE INVENTION

30 As discussed above, in one embodiment, the above-described objects of the present invention have been met by a non-viral:DNA complex comprising DNA encoding an anti-angiogenic peptide and DNA encoding a tumor suppressor protein.

The particular non-viral carrier (liposomes-neutral or non-cationic, see below) , polyethylimine (Fluka), polylysine (Sigma), polyhistidine (Sigma) , Superfect (Qiagen), are not critical to the present invention although cationic liposomes are preferable carriers. Examples of cationic lipids which can be employed in the present invention include include 1,2-
35 dioleoyl-sn-glycero-3-ethylphosphocholine (Avanti, Birmingham, AL), 1,2-dimyristoyl-sn-glycero-3-ethylphosphocholine (Avanti, Birmingham, AL), and (2,3-diol-eyloxy)propyl-N,N,N-trimethylammonium chloride (DOTMA) (Syntex Corp., Palo Alto, CA).

The cationic lipids are preferably used in a mixture with dioleoylphosphatidylethanolamine (DOPE) (Avanti, Birmingham, AL). In the present invention, the amount of cationic lipid present in the mixture is generally in the range of from 100%
40 to 40% (w/w), preferably about 50% (w/w); and the amount of DOPE present in the mixture is generally in the range of from 0% to 60% (w/w), preferably about 50% (w/w); and the amount of pegylated lipid (1,2 -diacyl-sn-glycero-3-phosphoethanolamine-N-[Poly(ethylene glycol) 2000] present in the mixture is generally in the range of from 0% to 10% (w/w), preferably about 1% (w/w).

45 The particular ligand will be dependent on the the tumor/peritumoral targeted. Examples of targets on tumors include Her2 (breast), CEA (colon), ferritin receptor (breast, lung, and ovary), and the tumor vasculature (α vB3 integrins or tissue factor). Antibodies directed toward Her2, CEA, and the tumor's vasculature will be coupled to 1% of the pegylated lipid hydroxyl group of the pegylated lipid with a water soluble carbodiimide (1-ethyl-3 (3-dimethylamino-propyl) carbodiimide), and purified over a sepharose CL-6B column. Similarly, ligands to the tumor (ferritin) and/or the vas-
50 culature (the peptide, RGD) are covalently attached to the hydroxyl the pegylated lipids.

The particular tumor suppressor gene employed is not critical to the present invention. Examples of such tumor suppressor genes include the p53 gene, the p21 gene (El-Deiry et al, *supra*; and Harper, *supra*), and the rb gene (Bookstein et al, *supra*). The p53 gene is the preferred tumor suppressor gene employed in the present invention.

55 The particular anti-angiogenic peptide encoded by the DNA is not critical to the present invention. Examples of said peptides include a fragment of thrombospondin I (TSP1) having the following amino acid sequence (the amino acid sequences that are known to be anti-angiogenic are underlined):

MTEENKELANELRRPPLCYHNGVQYRNNEEWTVDSCTECHCQNSVTICKKVSCP
 IMPCSNATVPDGECCPRCWPSDSADDGWSPWSEWTSCSTSCGNGIQQRGRSCDS
 5 LNNRCEGSSVQTRTCHIQECDKRFKQDGGWSHWSPWSSCSVTCGDGVITRITLC
 NSPSPQMNGKPCEGEARETKACKKDACPINGGWGPWSPWDICSVTCGGGVQKRS
 10 RL (SEQ ID NO:1),

which is encoded by the following DNA sequence (nucleotides 1013-1650 of the TSP1 gene; the underlined sequences encode the anti-angiogenic peptides; the bold TAA is the stop codon):

15 ATGACTGAAGAGAACAAAGAGTTGGCCAATGAGCTGAGGCGGCTCCCT
ATGCTATCACAAACGGAGTTCACTACAGAAATAACGAGGAATGGACTGTTG
 ATAGCTGCACTGAGTGTCACTGTCAGAACTCAGTTACCATCTGCAAAAAG
 20 GTGTCCTGCCCCATCATGCCCTGCTCCAATGCCACAGTTCTGATGGAGA
 ATGCTGTCCTCGCTGTTGGCCAGCGACTCTGCGGACGATGGCTGGTCTC
 CATGGTCCGAGTGGACCTCCTGTTCTACGAGCTGTGGCAATGGAATTCAG
 25 CAGCGCGGCCGCTCCTGCGATAGCCTCAACAACCGATGTGAGGGCTCCTC
 GGTCCAGACACGGACCTGCCACATTCAGGAGTGTGACAAAAGATTAAAC
 AGGATGGTGGCTGGAGCCACTGGTCCCGTGGTCATCTTGTCTGTGACA
 30 TGTGGTGATGGTGTGATCACAAGGATCCGGCTCTGCAACTCTCCAGCCC
 CCAGATGAATGGGAAACCCTGTGAAGGCGAAGCGCGGGAGACCAAAGCCT
 GCAAGAAAGACGCCTGCCCCATCAATGGAGGCTGGGGTCCTTGGTCACCA
 35 TGGGACATCTGTTCTGTACCTGTGGAGGAGGGGTACAGAAACGTAGTCT
 CTCTAA (SEQ ID NO:2);

a concatamer of TSP1 having the following amino acid sequence (the intervening sequence is underlined):

40 MTEENKELANELRRPPLCYHNGVQYRNNEEWTVDSCTECHCQNSVTICKKVSCP
 IMPCSNATVPDGECCPRCWPSDSADDGWSPWSEWTSCSTSCGNGIQQRGRSCDS
 45 LNNRCEGSSVQTRTCHIQECDKRFKQDGGWSHWSPWSSCSVTCGDGVITRITLC
 NSPSPQMNGKPCEGEARETKACKKDACPINGGWGPWSPWDICSVTCGGGVQKRS

50

55

RLCVHSRMTEENKELANELRRPPLCYHNGVQYRNNEEWTVDSCTECHCONSVTI
 CKKVSCPIMPCSNATVPDGECCPRCWPSDSADDGWSPWSEWTSCSTSCGNGIQQ
 5 RGRSCDSLNNRCEGSSVQTRTCHIIECDKRFKQDGGWSHWSPWSSCSVTGCDGV
 ITRITLCNSPSPQMNGKPCEGEARETKACKKDACPINGGWGPWSPWDICSVTCG
 GGVQKRSRL (SEQ ID NO:3),

10 which is encoded by the following DNA sequence (the intervening sequence is underlined):

ATGACTGAAGAGAACAAAGAGTTGGCCAATGAGCTGAGGCGGCCTCCCC
 15 TATGCTATCACACGGAGTTCAGTACAGAAATAACGAGGAATGGACTGTT
 GATAGCTGCACTGAGTGTCACTGTCAAGACTCAGTTACCATCTGCAAAAA
 GGTGTCTTGGCCCATCATGCCCTGCTCCAATGCCACAGTTCTTGATGGAG
 20 AATGCTGTCTCGCTGTTGGCCAGCGACTCTGCGGACGATGGCTGGTCT
 CCATGGTCCGAGTGGACCTCCTGTTCTACGAGCTGTGGCAATGGAATTCA
 GCAGCGCGGCCGCTCCTGCGATAGCCTCAACAACCGATGTGAGGGCTCCT
 25 CGGTCCAGACACGGACCTGCCACATTCAGGAGTGTGACAAAAGATTTAA
 CAGGATGGTGGCTGGAGCCACTGGTCCCCGTGGTCATCTTGTTCTGTGAC
 ATGTGGTGATGGTGTGATCACAAGGATCCGGCTCTGCAACTCTCCCAGCC
 CCCAGATGAATGGGAAACCCTGTGAAGGCGAAGCGCGGGAGACCAAAGCC
 30 TGCAAGAAAGACGCCTGCCCCATCAATGGAGGCTGGGGTCTTGGTCACC
 ATGGGACATCTGTTCTGTACCTGTGGAGGAGGGGTACAGAAACGTAGTC
 GTCTCTGCGTCGACTCTAGAATGACTGAAGAGAACAAAGAGTTGGCCAA
 35 TGAGCTGAGGCGGCCTCCCCATGCTATCACACGGAGTTCAGTACAGAA
 ATAACGAGGAATGGACTGTTGATAGCTGCACTGAGTGTCACTGTCAAGAC
 TCAGTTACCATCTGCAAAAAGGTGTCCTGCCCCATCATGCCCTGCTCCAA
 40 TGCCACAGTTCTTGATGGAGAATGCTGTCTCGCTGTTGGCCAGCGACT
 CTGCGGACGATGGCTGGTCTCCATGGTCCGAGTGGACCTCCTGTTCTACG
 AGCTGTGGCAATGGAATTCAGCAGCGCGGCCGCTCCTGCGATAGCCTCAA
 45 CAACCGATGTGAGGGCTCCTCGGTCCAGACACGGACCTGCCACATTCAGG
 AGTGTGACAAAAGATTTAAACAGGATGGTGGCTGGAGCCACTGGTCCCCG
 TGGTCATCTTGTTCTGTGACATGTGGTGATGGTGTGATCACAAGGATCCG
 50 GCTCTGCAACTCTCCCAGCCCCAGATGAATGGGAAACCCTGTGAAGGCG
 AAGCGCGGGAGACCAAAGCCTGCAAGAAAGACGCCTGCCCCATCAATGGA
 GGCTGGGGTCTTGGTCACCATGGGACATCTGTTCTGTACCTGTGGAGG
 55 AGGGGTACAGAAACGTAGTCGTCTCTAA (SEQ ID NO:4);

laminin peptide having the following amino acid sequence: MYIGSR (SEQ ID NO:5), which is encoded by the following

DNA sequence (the *Sa*II sites are underlined, and the stop codon is in bold):

GTCGACATGTATATTGGTTCTCGTTAAGTCGAC (SEQ ID NO:6);

a concatamer of the laminin sequence having the following amino acid sequence (the intervening sequences are underlined):

- 5 MYIGSRGKSYIGSRGKSYIGSRGKS (SEQ ID NO:7), which is encoded by the following DNA sequence (the *Sa*II sites are underlined, and the intervening sequences are in bold):

GTCGACATGTATATTGGTTCTCGTGGTAAAAGATATATTGGTTCTCGTGGTAAA

- 10 AGATATATTGGTTCTCGTGGTAAAAGATAAGTCGACC (SEQ ID NO:8);

a peptide from platelet factor-4 having the following amino acid sequence:

MLYKKIIKKLLES (SEQ ID NO:9), which is encoded by the following DNA sequence (the *Sa*II sites are underlined):

- 15 GTCGACATGCTTTATAAGAAGATCATCAAGAAGCTTCTTGAGAGTTAAGTCGAC (SEQ ID NO:10);

a concatamer of the platelet factor-4 peptide having the following amino acid sequence (the intervening sequences are underlined):

MLYKKIIKKLLESGKSLYKKIIKKLLESGKSLYKKIIKKLLESGK (SEQ ID NO:11), which is encoded by the following DNA sequence (the *Sa*II sites are underlined, and the intervening sequences are in bold):

- 20 GTCGACATGCTTTATAAGAAGATCATCAAGAAGCTTCTTGAGAGTGGTAAAAGA
CTTTATAAGAAGATCATCAAGAAGCTTCTTGAGAGTGGTAAAAGACTTTATAAG

- 25 AAGATCATCAAGAAGCTTCTTGAGAGTGGTAAAAGATAAGTCGAC (SEQ ID NO:12);

somatostatin inhibitor having the following amino acid sequence: MFCYWKCW (SEQ ID NO:13), which is encoded by the following DNA sequence (the *Sa*II sites are underlined):

- 30 GTCGACATGTTCTTGTATTGGAAGGGATTGTGGTAAAGTCGAC (SEQ ID NO:14);

a concatamer of somatostatin inhibitor having the following amino acid sequence (the intervening sequences are underlined):

- 35 MFCYWKCWGKSFCYWKCWGKSFCYWKCWGKS (SEQ ID NO:15), which is encoded by the following DNA sequence (the *Sa*II sites are underlined, and the intervening sequences are in bold):

- GTCGACATGTTCTTGTATTGGAAGGGATTGTGGGGTAAAAGATTCTTGTATTGG
 40 AAGGGATTGTGGGGTAAAAGATTCTTGTATTGGAAGGGATTGTGGGGTAAAAGA
TAAGTCGAC (SEQ ID NO:16);

fibronectin inhibitor having the following amino acid sequence: MGRGD (SEQ ID NO:17), which is encoded by the following DNA sequence (the *Sa*II sites are underlined):

- 45 GTCGACATGTCTTTGTCTTGAAGACTTTGACTTAAGTCGAC (SEQ ID NO:18);

a concatamer of fibronectin inhibitor having the following amino acid sequence (the intervening sequences are underlined):

- 50 MGRGDGKSGRGDGKSGRGDGKS (SEQ ID NO:19); which is encoded by the following DNA sequence (the *Sa*II sites are underlined, and the intervening sequences are in bold):

GTCGACATGGGTCGTGGTGATGGTAAAAGAGGTCGTGGTGATGGTAAAAGA

GGTCGTGGTGATGGTAAAAGATAAGTCGAC (SEQ ID NO:20);

5

angiostatin having the following amino acid sequence:

MVYLSECKTGIGNGYRGTMSTRKSGVACQKWGATFPHVPNYSPSTHPNEGLEEN

YCRNPDNDEQGPWCYTTPDKRYDYCNIPCEEEECMYCSGEKYEGKISKTMMSG

10

LDCQAWDSQSPHAHGYIPAKFP SKNLKMNYCHNPDGEPRPWCFTTDP TKRWEYC

DIPRCTTPPPPPSPTYQCLKGRGENYRGTVSVTVSGKTCQRWSEQTPHRHNRTF

ENFPCKNLEENYCRNPDGETAPWCYT TDSQLRWEYCEIPSCCESSASPDQSDSSV

15

PPEEQTPVVQECYQSDGQSYRGTSSTTITGKKCQSEQTPHR

(SEQ ID NO:21),

20 which is encoded by the following DNA sequence (the *Sall* sites are underlined):

GTCGACATGGTGTATCTGTCAGAATGTAAGACCGGCATCGGCAACGGC

TACAGAGGAACCATGTCCAGGACAAAGAGTGGTGTGCGCTGTCAAAG

25

TGGGGTGCCACGTTCCCCACGTACCCAACTACTCTCCCAGTACACAT

CCCAATGAGGGACTAGAAGAGAACTACTGTAGGAACCCAGACAATGAT

GAACAAGGGCCTTGGTGCTACACTACAGATCCGGACAAGAGATATGAC

30

TACTGCAACATTCTGAATGTGAAGAGGAATGCATGTACTGCAGTGGA

GAAAAGTATGAGGGCAAATCTCCAAGACCATGTCTGGACTTGACTGC

CAGGCCTGGGATTCTCAGAGCCACATGCTCATGGATACATCCCTGCC

35

AAATTTCCAAGCAAGAACCTGAAGATGAATTATTGCCACAACCCTGAC

GGGGAGCCAAGGCCCTGGTGCTTCACAACAGACCCCAACAAACGCTGG

GAATACTGTGACATCCCCGCTGCACAACACCCCGCCCCACCCAGC

40

CCAACCTACCAATGTCTGAAAGGAAGAGGTGAAAATTACCGAGGGACC

GTGTCTGTCACCGTGTCTGGGAAAACCTGTCAGCGCTGGAGTGAGCAA

ACCCCTCATAGGTGA GTCGAC (SEQ ID NO:22),

45

a concatamer of angiostatin having the following amino acid sequence (the intervening sequences are underlined):

50

55

MVYLSECKTGIGNGYRGTMSTRKSGVACQKWGATFPHVPNYSPSTHPNEGLE
 ENYCRNPDNDEQGPWCYTTPDPDKRYDYCNIPCEEEECMYCSGEKEYEGKISKTMS
 5 GLDCQAWDSQSPHAHGYIPAKFPSKNLKMNYCHNPDGEPRPWCFTTDPDKRWEY
 CDIPRCTTPPPPPSPTYQCLKGRGENYRGTVSVTVSGKTCQRWSEQTPHRHNRT
 PENFPCKNLEENYCRNPDGETAPWCYTTPDSQLRWEYCEIPSCCESSASPDQSDSS
 10 VPPEEQTPVVQECYQSDGQSYRGTSSTTITGKKCQSEQTPHRGKSMVYLSECKT
 GIGNGYRGTMSTRKSGVACQKWGATFPHVPNYSPSTHPNEGLEENYCRNPDNDE
 QGPWCYTTPDPDKRYDYCNIPCEEEECMYCSGEKEYEGKISKTMSGLDCQAWDSQS
 15 PHAHGYIPAKFPSKNLKMNYCHNPDGEPRPWCFTTDPDKRWEYCDIPRCTTPPP
 PPSPTYQCLKGRGENYRGTVSVTVSGKTCQRWSEQTPHRHNRTPENFPCKNLEE
 NYCRNPDGETAPWCYTTPDSQLRWEYCEIPSCCESSASPDQSDSSVPPEEQTPVVQ
 20 ECYQSDGQSYRGTSSTTITGKKCQSEQTPHR (SEQ ID No :23),

which is encoded by the following DNA sequence (the *Sal*I sites are underlined, and the intervening sequences are in bold):

25 GTCGACATGGTGTATCTGTCAGAAATGTAAGACCGGCATCGGCAACGGCTACA
 GAGGAACCATGTCCAGGACAAAGAGTGGTGTTCCTGTCAAAAGTGGGGTGC
 30 CACGTTCCCCCACGTACCCAACTACTCTCCAGTACACATCCCAATGAGGGA
 CTAGAAGAGAACTACTGTAGGAACCCAGACAATGATGAACAAGGGCCTTGGT

GCTACACTACAGATCCGGACAAGAGATATGACTACTGCAACATTCCTGAATG
 TGAAGAGGAATGCATGTACTGCAGTGGAGAAAAGTATGAGGGCAAATCTCC
 5 AAGACCATGTCTGGACTTGACTGCCAGGCCTGGGATTCTCAGAGCCCACATG
 CTCATGGATACATCCCTGCCAAATTTCCAAGCAAGAACCTGAAGATGAATTA
 TTGCCACAACCCTGACGGGGAGCCAAGGCCCTGGTGCTTCACAACAGACCCC
 10 ACCAAACGCTGGGAATACTGTGACATCCCCCGCTGCACAACACCCCCGCCCC
 CACCCAGCCCAACCTACCAATGTCTGAAAGGAAGAGGTGAAAATTACCGAGG
 GACCGTGTCTGTACCGTGTCTGGGAAAACCTGTCAGCGCTGGAGTGAGCAA
 15 ACCCCTCATAGGGGTAAAAGAATGGTGTATCTGTGAGAATGTAAGACCGGCA
 TCGGCAACGGCTACAGAGGAACCATGTCCAGGACAAAGAGTGGTGTTCCTG
 TCAAAAGTGGGGTGCCACGTTCCCCACGTACCCAATACTCTCCAGTACA
 20 CATCCCAATGAGGGACTAGAAGAGAACTACTGTAGGAACCCAGACAATGATG
 AACAAGGGCCTTGGTGCTACACTACAGATCCGGACAAGAGATATGACTACTG
 CAACATTCCTGAATGTGAAGAGGAATGCATGTACTGCAGTGGAGAAAAGTAT
 25 GAGGGCAAATCTCCAAGACCATGTCTGGACTTGACTGCCAGGCCTGGGATT
 CTCAGAGCCCACATGCTCATGGATACATCCCTGCCAAATTTCCAAGCAAGAA
 CCTGAAGATGAATATTGCCACAACCCTGACGGGGAGCCAAGGCCCTGGTGCT
 30 TCACAACAGACCCCAACAAACGCTGGGAATACTGTGACATCCCCCGCTGCACAA
 CACCCCCGCCCCACCCAGCCCAACCTACCAATGTCTGAAAGGAAGAGGTGAAA
 ATTACCGAGGGACCGTGTCTGTACCGTGTCTGGGAAAACCTGTCAGCGCTGGA
 35 GTGAGCAA ACCCCTCATAGGTGA GTCGAC (SEQ ID NO:24);

prolactin having the following amino acid sequence:

40 MLPICPGGAARCQVTLRDLFDRAVVLSHYIHNLSSEMFSEFDKRYTHGRGFI
 TKAINSCHTSSLATPEDKEQAQOMNQKDFLSLIVSILRSWNEPLYHLVTEVR
 GMQEAPEAILSKAVEIEEQTK (SEQ ID NO:25),

45 which is encoded by the following DNA sequence:
 50
 55

ATGTTGCCCATCTGTCCCGCGGGGCTGCCCGATGCCAGGTGACCCTTCGAG
 ACCTGTTTGACCGCGCCGTCGTCTGTCCCACTACATCCATAACCTCTCCTC
 5 AGAAATGTTTCAGCGAATTCGATAAACGGTATACCCATGGCCGGGGGTTTCATT
 ACCAAGGCCATCAACAGCTGCCACACTTCTTCCCTTGCCACCCCCGAAGACA
 AGGAGCAAGCCCAACAGATGAATCAAAAAGACTTTCTGAGCCTGATAGTCAG
 10 CATATTGCGATCCTGGAATGAGCCTCTGTATCATCTGGTCACGGAAGTACGT
 GGTATGCAAGAAGCCCCGGAGGCTATCCTATCCAAAGCTGTAGAGATTGAGG
 AGCAAACCTAA (SEQ ID NO:26);

15 and
 a concatamer of prolactin having the following amino acid sequence (the intervening sequences are underlined):

20 MLPICPGGAARCQVTLRDLFDRAVVLSHYIHNLSSEMFSEFDKRYTHGRGFITK
 AINSCHTSSLATPEDKEQAQOMNQKDFLSLIVSILRSWNEPLYHLVTEVRGMQE
 APEAILSKA~~VEIEEQTKGK~~SMLPICPGGAARCQVTLRDLFDRAVVLSHYIHNLS
 25 SEMFSEFDKRYTHGRGFITKAINSCHTSSLATPEDKEQAQOMNQKDFLSLIVSI
 LRSWNEPLYHLVTEVRGMQEAP~~EAILS~~KAVEIEEQTK (SEQ ID NO:27),

30 which is encoded by the following DNA sequence (the intervening sequences are in bold):

ATGTTGCCCATCTGTCCCGCGGGGCTGCCCGATGCCAGGTGACCCTTCGAG
 ACCTGTTTGACCGCGCCGTCGTCTGTCCCACTACATCCATAACCTCTCCTC
 35 AGAAATGTTTCAGCGAATTCGATAAACGGTATACCCATGGCCGGGGGTTTCATT
 ACCAAGGCCATCAACAGCTGCCACACTTCTTCCCTTGCCACCCCCGAAGACA
 AGGAGCAAGCCCAACAGATGAATCAAAAAGACTTTCTGAGCCTGATAGTCAG
 40 CATATTGCGATCCTGGAATGAGCCTCTGTATCCTGGTCACGGAAGTACGT
 GGTATGCAAGAAGCCCCGGAGGCTATCCTATCCAAAGCTGTAGAGATTGAGG
 AGCAAACCGGTAAAGAAATGTTGCCCATCTGTCCCGCGGGGCTGCCCGATGCC
 45 AGGTGACCCTTCGAGACCTGTTTGACCGCGCCGTCGTCTGTCCCACTACATCC
 ATAACCTCTCCTCAGAAATGTTTCAGCGAATTCGATAAACGGTATACCCATGGCC
 GGGGGTTTCATTACCAAGGCCATCAACAGCTGCCACACTTCTTCCCTTGCCACCC
 50 CCGAAGACAAGGAGCAAGCCCAACAGATGAATCAAAAAGACTTTCTGAGCCTGA
 TAGTCAGCATATTGCGATCCTGGAATGAGCCTCTGTATCATCTGGTCACGGAAG
 TACGTGGTATGCAAGAAGCCCCGGAGGCTATCCTATCCAAAGCTGTAGAGATTG
 55 AGGAGCAAACCTAA (SEQ ID NO:28)

Increase efficacy will occur with concatamers of the anti-angiogenic genes. This will increase the anti-angiogenic dosage level without changing the amount of vector necessary to deliver these genes. Similar to concatamers, a plasmid with two or more promoters, a plasmid with the IRES sequence (internal ribosomal entry site) between two sequences, and an antiangiogenic peptide with a secretory sequence will increase the delivery of genes to the therapeutic target without markedly increasing the DNA concentration. With regards to the concatamers, the concatamers can extend up to approximately 4400 bases in length (the coding region of a large protein), and the number of concatamers possible will depend on the number of bases of a single anti-angiogenic unit.

For fibronectin, the range of concatamers would be about 2 to 66. Although, the maximum number of anti-angiogenic units for the TSPf is about 6, one can increase this concatameric number by deleting the sequences that do not have any anti-angiogenic effects, such as shown below:

ATG (CTGAGGCGGCCTCCCCTATGCTATCACAAACGGAGTTCAGTACAGAAATAA
CGGTAAAAGATCCCCGTGGTCATCTTGTCTGTGACATGTGGTGATGGTGTGAT
GGTAAAAGAAGTGGTACCCTGTAGACAAGACAGTGGACACCTCCTCCCCAT)_nT
AA (SEQ ID NO:29),

where the corresponding amino acid sequence is:

M(LRRPPLCYHNGVQYRNNEEWTVDSGKSSPWSSCSVTCGDGVITRIGKSSPWD
ICSVTCGGGV)_n (SEQ ID NO:30),

and wherein n is an integer of from 2 to 24. In a similar manner, the concatameric number of the platelet factor-4 peptide, somatostatin inhibitor, angiostatin, and prolactin can be increased.

Since more than one anti-angiogenic pathway exists, concatamers consisting of two or more types of inhibitor are believed to be more effective than the homogenous concatamers. For example, heterogeneous concatamers of TSP1 and the fibronectin inhibitors can be inserted into the same vector. An example of a heterogeneous concatamer useful in the present invention is as follows:

ATG (CTGAGGCGGCCTCCCCTATGCTATCACAAACGGAGTTCAGTACAGAAATAA
CGGTAAAAGATCCCCGTGGTCATCTTGTCTGTGACATGTGGTGATGGTGTGAT
GGTAAAAGAAGTGGTACCCTGTAGACAAGACAGTGGACACCTCCTCCCCAT)_x(
TATATTGGTTCTCGTGGTAAAAGA)_yTAA (SEQ ID NO:31).

The first parenthetical represents the nucleotide sequence of TSPf, whereas the second parenthetical represents the anti-angiogenic fragment isolated from fibronectin, wherein x and y represent the number of repeats of TSP1 and fibronectin, respectively. Again, the number of bases delineated by the summation of x + y will generally not exceed 4400 bases.

The above heterogeneous concatamers need not be limited to only anti-angiogenic peptides. For example, the protein angiostatin or the large polypeptide fragment of prolactin can be modified with the above-mentioned genes which encode anti-angiogenic peptides. Again, the concatameric number will vary depending on the number of nucleotide bases of the unit angiogenic inhibitor. In this concatamer of large and small anti-angiogenic inhibitors, the ratio of large to small inhibitors is 0.1 to 0.9, preferably 1:1.

A translational start signal Met, has been included in all of the above peptides; and a transcriptional stop codon (TAA) has been included in all of the above DNA sequences.

The *Sall* sites indicated in the above-sequences are a useful cloning tool for insertion of the DNA into BAP vector, which is known to be useful in expressing proteins efficiently *in vivo* from the β -actin promoter (Ray et al, *Genes Dev.*, 5:2265-2273 (1991)). Other restriction sites can be incorporated into the DNA for cloning into other vectors.

Other useful vectors for gene therapy which can be employed in the present invention include plasmids with a simian viral promoter, e.g., pZeoSV (Invitrogen); or the CMV promoter, e.g., pcDNA3, pRc/CMV or pcDNA1 (Invitrogen). Plasmids with a CMV promoter may contain an intron 5' of the multiple cloning site (Zhu et al, *supra*). Plasmids contain-

ing the BGH terminator instead of the viral SV40 polyA terminator, e.g., pcDNA3, pRc/CMV, pRc/RSV (Norman et al, IBC's 5th Annual Meeting (1995); and Invitrogen vectors), can also be employed in the present invention so as to increase the expression of the tumor suppressor gene and the anti-angiogenic peptide in cells. As stated previously, a vector containing two or more promoters will greatly enhance the therapeutic efficacy. Vectors containing the IRES sequence which allows the translation of two different coding genes to occur from one mRNA transcript will also significantly increase the efficacy of the therapy.

Expression of the DNA encoding the tumor suppressor protein and the DNA encoding the anti-angiogenic peptide can be achieved using a variety of promoters, and the particular promoter employed is not critical to the present invention. For example, the promoter can be a generalized promoter, such as the β -actin promoter, a simian viral promoter, or the CMV promoter, or a tissue specific promoter, such as the α -fetal protein promoter which is specific for liver (Kaneko et al, *Cancer Res.*, 55:5283-5287 (1995), the tyrosinase promoter which is specific for melanoma cells (Hughes et al, *Cancer Res.*, 55:3339-3345 (1995); or the enolase promoter which is specific for neurons (Andersen et al, *Cell. Molec. Neurobiol.*, 13:503-515 (1993)).

The particular amount of DNA included in the cationic liposomes of the present invention is not critical. Generally, the amount of total DNA is in the range of about 0.005 to 0.32 $\mu\text{g}/\mu\text{M}$ of liposome, preferably 0.045 to 0.08 $\mu\text{g}/\mu\text{M}$ of liposome.

The DNA encoding a tumor suppressor gene is generally present in an amount of from 0.0025 to 0.16 $\mu\text{g}/\mu\text{M}$ of liposome, preferably 0.028 to 0.04 $\mu\text{g}/\mu\text{M}$ of liposome. The DNA encoding an anti-angiogenic peptide is generally present in an amount of from 0.0025 to 0.16 $\mu\text{g}/\mu\text{M}$ of liposome, preferably 0.028 to 0.04 $\mu\text{g}/\mu\text{M}$ of liposome.

The mole ratio of the DNA encoding the tumor suppressor gene to the DNA encoding the anti-angiogenic peptide is not critical to the present invention. Generally, the mole ratio is between 1:5 to 5:1, preferably about 1 to 1.

The DNA encoding the tumor suppressor gene and the anti-angiogenic peptide may be contained on the same vector, or on separate vectors.

Cationic liposomes are prepared similarly to other liposomes. In brief, the cationic lipid with/or without DOPE are dissolved in a solvent, e.g., chloroform. The lipids are then dried in a round bottom flask overnight on a rotary evaporator. The resulting lipids are then hydrated with sterile water over a 1 hr period so form large multilamellar vesicle liposomes. To decrease the size of the liposomes, one may sonicate or pass the liposomes back and forth through a polycarbonate membrane. The DNA is then added to a solution containing the liposomes after their formation.

In another embodiment, the above-described objects of the present invention have been met by a method for inhibiting tumor growth in a subject comprising administering to a tumor-bearing subject a cationic liposome:DNA complex comprising DNA encoding a tumor suppressor gene and DNA encoding an anti-angiogenic peptide.

In a preferred embodiment the cationic liposome:DNA complex additionally comprises DNA encoding a tumor suppressor protein.

In a further preferred embodiment the cationic polymer:DNA complex additionally comprises DNA encoding a tumor suppressor protein. In a further preferred embodiment the cationic liposomes in the cationic liposome:DNA complex are comprised of one cationic lipid (i.e., 1,2-dioleoyl-sn-glycero-3-ethyl-phosphocholine, 1,2-dimyristoyl-sn-glycero-3-ethylphosphocholine, and 2,3-diol-eyloxy)propyl-N,N,N-tri-methyl-ammonium chloride) and may also be comprised of polyethylimine glycol (i.e., a pegylated lipid-1,2-diacy-sn-glycero-3-phospho-ethanolamine-N-Poly(ethylene glycol) 2000) and a targeted ligand (the amino acid peptide RGD, ferritin, or antibodies targeted toward to HER2).

In a further preferred embodiment of the invention said cationic liposomes in said complexes are comprised of one cationic polymer polyethylimine, polylysine, polyhisitidine, and Superfect) and may also be comprised of polyethylene glycol and a targeted ligand (the amino acid peptide RGD, ferritin, or antibodies targeted toward to HER2, CEA).

In a further preferred embodiment said tumor suppressor protein in said complexes is selected from the group consisting of the p53, the p21 and the rb.

In a further preferred embodiment said tumor suppressor protein in the said complexes is p53.

In a further preferred embodiment of the invention said anti-angiogenic peptide in the said complexes is selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, and SEQ ID NO:30.

In a further preferred embodiment the said DNA encoding an anti-angiogenic peptide used in the said cationic liposome:DNA complex is present in an amount of from 0.0025 to 0.16 $\mu\text{g}/\mu\text{M}$ of liposome.

In a further preferred embodiment the said DNA encoding an anti-angiogenic peptide in the said cationic polymer:DNA complex is present in an amount of from 0.016 to 0.33 $\mu\text{g}/\mu\text{g}$ of polymer.

In a further preferred embodiment the said DNA encoding a tumor suppressor protein in the said complexes is present in an amount of from 0.0025 to 0.16 $\mu\text{g}/\mu\text{M}$.

In a further preferred embodiment of the present invention said DNA encoding a tumor suppressor protein in the said complexes is present in an amount of from 0.016 to 0.33 $\mu\text{g}/\mu\text{M}$.

A further embodiment of the present invention is the provision of the use of a cationic polymer:DNA complex com-

prising DNA encoding an anti-angiogenic peptide for the production of a medicament for inhibiting tumor growth in a subject which preferably comprises administering the same to a tumor-bearing subject.

In a further preferred embodiment of the present invention the said complex in the said use additionally comprises DNA encoding a tumor suppressor protein.

5 In a further preferred embodiment of the present invention the said cationic liposome in the said use is (i.e. 1,2-dioleoyl-sn-glycero-3-ethylphosphocholine, 1,2-dimyristoyl-sn-glycero-3-ethylphosphocholine, and 2,3-dioleoylpropyl-N,N,N-trimethylammonium chloride and may also be comprised of polyethylene glycol (i.e., a pegylated lipid-1,2-diacyl-sn-glycero-3-phosphoethanolamine-N-[Poly-(ethylene glycol) 2000] and a targeted ligand (the amino acid peptide RGD, ferritin, or antibodies targeted toward to HER2).

10 In a further preferred embodiment of the present invention the said cationic polymer in the said use is (i.e. (polyethylimine Polycat57, polylysine, polyhistidine, and Superfect) and may also be comprised of polyethylene glycol and a targeted ligand (the amino acid peptide RGD, ferritin, or antibodies targeted toward to HER2).

In a further preferred embodiment of the present invention the said tumor suppressor protein used in the said cationic polymer is selected from the group consisting of p53, the p21 and the rb.

15 In a further preferred embodiment of the present invention the said tumor suppressor protein used in the said cationic polymer is p53.

In a further preferred embodiment the said anti-angiogenic peptide used for the said cationic polymer is selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, and SEQ ID NO:30.

20 In a further preferred embodiment of the present invention the said DNA encoding an anti-angiogenic peptide used in the cationic complex is present in an amount from 0.0025 to 0.16 $\mu\text{g}/\text{pM}$ of liposome or 0.016 to 0.33 $\mu\text{g}/\mu\text{g}$ of polymer.

25 In a further preferred embodiment of the present invention the said DNA encoding a tumor suppressor protein used in the said complex is present in an amount of from 0.0025 to 0.16 $\mu\text{g}/\text{pM}$ of liposome or 0.016 to 0.33 $\mu\text{g}/\mu\text{g}$ of polymer.

In a further embodiment of the present invention the said cationic polymer:DNA complex comprising a plasmid with one or more promoters expressing either the same gene product (SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, and SEQ ID NO:30) or combinations of these gene products.

30 In a further embodiment of the present invention the cationic liposome:DNA complex comprising a plasmid with an intervening internal ribosomal entry site sequence between two genes (SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:22, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, and SEQ ID NO:29) that are either the same or different.

35 In a further preferred embodiment the said anti-angiogenic protein in the said cationic liposome:DNA complex or the said cationic polymer:DNA complex is secretory from the group consisting of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, and SEQ ID NO:30.

40 The particular type of tumor which can be treated in the present invention is not critical thereto. Examples of tumors which can be treated in accordance with the present invention include solid tumors, e.g., lung, colon, brain, breast and melanoma tumors. All of these tumors are very dependent on blood supply to sustain their growth.

The particular mode of administering the cationic liposome:DNA complex of the present invention is not critical thereto. Examples of such modes include intravenous, subcutaneous and intratumoral injections. Intravenous injection is the preferred administration mode since there is better distribution to the developing blood vessels of the tumor.

45 The amount of cationic liposome:DNA complex to be administered will vary depending upon the age, weight, sex of the subject, as well as the tumor volume and rate of tumor growth in the subject. Generally, the amount to be administered will be about 5 to 60 μg , preferably about 9 to 16 μg .

The following examples are provided for illustrative purposes only, and are in no way intended to limit the scope of the present invention.

50 EXAMPLE 1

Production of DNA Vectors

A. TSPI Vector

55

The coding region of the TSPI gene is well-known (GB Accession code-X14787). The TSPI gene was inserted into the *Xba*I site of BAP vector (Ray et al, *supra*), so as to give rise to TSPI vector, wherein expression of the TSPI gene is controlled by the β -actin promoter.

More specifically, TSPI cDNA and Bluescript plasmid (Promega) were digested with *Hind*I and *Xba*I, and then the TSPI cDNA was ligated into Bluescript. Next, Bluescript containing the TSP cDNA and BAP vector were digested with *Sal*I and *Bam*HI, and TSPI cDNA inserted in the *Xba*I site of BAP vector. The correct orientation of the TSPI gene in BAP vector was confirmed by DNA sequencing.

B. TSPf Vector

TSPf vector is a vector containing a DNA fragment of the TSPI gene which has the two anti-angiogenic domains (nucleotides 992-1650) (Tolsma et al, *supra*), and a start codon and a stop codon.

The DNA fragment was prepared by PCR using thrombospondin I cDNA as template, and 100 pmols of each of the following primers 5'-TAGG**TCTAG**AATGACTGAAGAGAACAAAGAG-3' (SEQ ID NO:24); and 5'-ATGG**TCTAG**ATT-AGAGACGACTACGTTTCTG-3' (SEQ ID NO:25) so as to amplify nucleotides 1013 to 1650 of the TSPI gene. Both primers contain *Xba*I sites (underlined), the first primer contains an ATG start codon (in bold), and the second primer contains a TTA stop codon (in bold).

The resulting 638 base pair fragment of the TSPI gene (hereinafter "TSPf") encodes the peptides that are known to be angiogenic inhibitors (Tolsma et al, *supra*).

After amplification, the DNA fragment was purified, digested with *Xba*I, and the digested fragment inserted into the *Xba*I site of BAP vector such that the expression of the TSPf gene was controlled by the β -actin promoter (Ray et al, *supra*; and Lesoon-Wood et al, *Human Gene Ther.*, 6:395-405 (1995)). The correct orientation of the fragment in BAP vector was verified by digestion with *Bam*HI, and confirmed by DNA sequencing.

C. p53 Vector

The coding sequence of the p53 gene was cut from plasmid p1SVhp53c62 (Zakut-Houri et al, *EMBO J.*, 4:1251-1255 (1985)) with *Xba*I, and inserted into the multiple cloning sites of pGEM3Z vector (Promega, Madison, WI). Digestion of the resulting vector with *Sal*I and *Bam*HI generated a 1900 bp fragment that was then inserted into the *Sal*I and *Bam*HI sites of BAP vector such that expression of the p53 gene was controlled by the β -actin promoter. The correct orientation of the p53 gene in BAP vector was confirmed by DNA sequencing.

EXAMPLE 2

Preparation of Cationic Liposome:DNA Complexes

A DOTMA:DOPE liposome mixture is known to efficiently transfect endothelial cells *in vitro* (Tilkins et al, *Focus*, 16:117-119 (1994)). Accordingly, liposome:DNA complexes were prepared using DOTMA:DOPE, in a 1:1 ratio, essentially as described by Debs et al, *J. Biol. Chem.*, 265:10189-10192 (1990). Similar liposomes preparations can be prepared by mixing, at a 1:1 ratio, DOPE with other cationic lipids, such as, 1,2-dioleoyl-sn-glycero-3-ethylphosphocholine, and 1,2-dimyristoyl-sn-glycero-3-ethylphosphocholine.

More specifically, a mixture of 400 nmols of the DOTMA and DOPE were dried overnight on a rotary evaporator. Then, the lipids were rehydrated with 1.5 ml of water for 2 hrs. Next, the milky liposome preparation was sonicated with a bath sonicator until clear. The resulting liposome preparation was then passed through a 50 nm polycarbonate filter between 15 to 20 times with a LipsoFast-Basic extruder (Avestin, Ottawa, On).

The DNA employed was either (1) empty BAP vector; (2) TSPI vector alone; (3) TSPf vector alone; (4) p53 vector alone; (5) p53 vector + TSPI vector; or (6) p53 vector + TSPf vector.

The DNA was prepared with the maxi Qiagen kits (Qiagen Inc., Chatsworth, Ca), and washed twice in 70% (v/v) ethanol. The DNA was then dialyzed against water for 24 hrs to removed any remaining salt.

About 400 pmols of the liposome preparation was gently mixed with between 18 to 35 μ g of total DNA in an Eppendorf tube. This amount in each eppendorf tube was sufficient for two injections. The same amount of DNA was injected in the combination therapies as in the single treatment regimens. For example, if 16 μ g of DNA in the combination therapy (8.0 μ g of p53 + 8.0 μ g of TSPf) was injected into each mouse of one group, then 16 μ g of p53 was injected into each mouse of a second group.

EXAMPLE 3

Anti-Angiogenic Effect of Cationic Liposome:DNA Complexes

The anti-angiogenic effects of the cationic liposome:DNA complexes obtained in Example 2 were evaluated in mice containing MDA-MB-435 breast cancer tumors (American Type Tissue Culture, Bethesda, MD), which are p53 defi-

cient.

More specifically, after administering the anesthetic, Metofane, to 126 female athymic nude mice (NCI), the mice were injected with 2.0×10^5 MDA-MB-435 tumor cells into the mammary fat pad using a stepper (Tridak) and a 27.5 g needle. Two weeks later, the mice whose tumors grew were divided into various treatment regimens, 18 mice per each regimen. The treatment regimens were as follows:

(1) untreated; (2) empty BAP vector; (3) TSP1 vector alone; (4) TSPf vector alone; (5) p53 vector alone; (6) p53 vector + TSP1 vector; and (7) p53 vector + TSPf vector. The mice received two intravenous injections, the first injection 14 days after the malignant cells had been implanted into the mice, and the second injection 24 days after the malignant cells had been implanted into the mice. The first injection consisted of 200 pmoles of the liposomes complexed with 16 μ g of total DNA. The second injection consisted of 200 pmoles of the liposomes complexed with 8.0 μ g of total DNA. The sizes of the tumors were measured 7 days after the second injection. The results are shown in Table 1 below.

TABLE 1

Anti-tumor Effects of TSP1 and TSPf	
Putative Anti-tumor Genes	Tumor Size (mm ³)
Untreated	113.5 \pm 6.41
BAP	102.9 \pm 6.83
TSP1	103.2 \pm 8.96
TSPf	89.4 \pm 11.06
p53	80.1 \pm 12.7*
p53 + TSP1	82.9 \pm 6.95*
p53 + TSPf	53.2 \pm 8.37**

* p53 or p53 + TSP1 vs. untreated, $p < 0.05$

** p53 + TSPf vs. untreated or BAP, $p < 0.01$

As shown in Table 1 above, the p53-treated group was found to be statistically different from the untreated group ($p < 0.05$) after 2 injections. However, the p53 treated group was not statistically different from the empty BAP vector group. This was similar to the results described by Lesoon-Wood et al, *Human Gene Ther.*, 6:395-406 (1995), in which p53 was not statistically different from the empty BAP vector group until after 5 injections.

However, p53 in combination with TSPf reduced tumors more effectively than p53 alone. That is, after just 2 injections of this combination therapy, there was a 35% further reduction in tumor growth compared to p53 alone. The combination group was statistically different from both the untreated and the empty BAP vector groups ($p < 0.01$). Although TSPf by itself was slightly less effective than p53, TSPf was, unexpectedly, substantially more effective than TSP1. In fact, the full length TSP1-treatment group had no more effect than either the empty vector or the untreated groups. This was unexpected for several reasons: 1) both the full length and the fragment of thrombospondin I contained the anti-angiogenic peptide 2) in a previous ex vivo study (Weinstat-Saslow et al, supra), full length thrombospondin I was effective in inhibiting tumor growth, and 3) full length thrombospondin I has a secretory sequence presumably so that the secreted protein can inhibit endothelial proliferation, whereas the thrombospondin I fragment does not contain a secretory sequence.

Regardless of whether there is a secretory sequence, one would predict prior to the present invention that the liposome:antiangiogenic gene would not be an effective antitumor therapy. As taught by Lesoon-Wood et al., the transfection efficiency of the tumor with cationic liposomes was very low. In fact, it could not be quantitated with a primer extension method. We know from the teaching of Weinstat-Saslow et al. that high levels of expressed TSP1 in 100% of the tumor cells reduces the tumor growth by only 60% in an ex vivo system. Extrapolating from these findings, a relatively high transfection efficiency of 20% with the liposome: antiangiogenic genes would have resulted in a marginal reduction ($20\%/100\% \times 60\% \text{ reduction} = 12\%$) of the tumor. This amount of tumor reduction would not have resulted in statistical differences with the liposome:antiangiogenic gene complexes. A transfection efficiency of the tumor above 10% would have easily been measurable with a variety of techniques including the primer extension method (used by Lesoon-Wood et al.). It has been determined that the transfection efficiency of the tumor is less than 5% with these cationic liposomes.

Hence, it was clearly unobvious that DNA encoding an anti-angiogenic peptide alone would be an effective anti-

tumor agent *in vivo*, based upon teachings that an anti-angiogenic peptide is an effective anti-tumor agent (Tolsma et al and Bouck et al), and based upon the teachings that DNA encoding a full-length anti-angiogenic protein is an effective anti-tumor agent *ex vivo* (Weinstat-Saslow et al).

A second experiment was carried out to determine whether the combination therapy of p53 and TSPf was effective at lower dosages, and to confirm that the combination of p53 and TSPf reduced the tumor size considerably more than p53 alone.

More specifically, 36 mice were injected with 2.0×10^5 MDA-MB-435 tumor cells into the mammary fat pad. Two weeks later, the mice whose tumors grew were divided into various treatment regimens, 18 mice per each regimen. The treatment regimens were as follows: (1) empty BAP vector; (2) p53 vector alone, and (3) p53 vector + TSPf vector. The mice were injected intravenously with 200 pmols of the liposomes complexed with 8.0 μ g of total DNA. Subsequently, the mice were treated in the same manner with 200 pmols of the liposomes complexed with 12 μ g of total DNA for the next 4 injections. Ten days elapsed between each injection. The sizes of the tumors were measured before each injection and 7 days after the last injection. The results are shown in Table 2 below:

TABLE 2

Anti-tumor Effects of p53 and TSPf	
Putative Anti-tumor Genes	Tumor Size (mm ³)
BAP	855 \pm 345
p53	616 \pm 142
p53 + TSPf	265 \pm 133*

* p53 + TSPf vs. BAP, $p < 0.02$

As shown in Table 2 above, the combination therapy with p53 and TSPf was statistically different from BAP, whereas the p53 alone treatment was not. This experiment confirmed that p53 and TSPf were more effective than p53 alone. Furthermore, a different dosage regimen, without an initial booster dose of 16 μ g of DNA as used in the experiment in Table 1, accentuated the difference between the combination treatment and the p53 alone treatments.

In table 1, the TSPf treatment group decreased the tumor more than empty vector or untreated groups. However, it was not statistically significant ($p = .07$). We repeated the experiment after injecting a higher dose of DNA and measured the different treatment groups tumors 10 days after the first treatment.

Table 3

Anti-tumor Effects of TSPf	
Putative Anti-tumor Genes	Tumor Size (mm ³)
Untreated	80.0 \pm 11.2
BAP	80.4 \pm 4.5
TSPf	50.7 \pm 4.8*

* TSPf vs. BAP, $p < 0.025$

At the higher dose of 19 μ g of DNA, the TSPf treatment group was statistically different from either the empty vector or the untreated groups.

In another experiment demonstrating the efficacy of antiangiogenic genes, various antiangiogenic genes were examined for their antitumor activity. After administering the anesthetic, Metofane, to 126 female athymic nude mice (NCI), the mice were injected with 3.0×10^5 MDA-MB-435 tumor cells into the mammary fat pad using a stepper (Tridak) and a 27.5 g needle. Two weeks later, the mice whose tumors grew were divided into various treatment regimens, 8 mice per each regimen. The treatment regimens were as follows: (1) BAP vector; (2) TSPf vector alone; (3) laminin peptide vector alone; and (4) angiostatin vector alone. The mice received 4 intravenous injections, the first injection was 10 days after the malignant cells had been implanted into the mice, and the remaining injections were thereafter 10 days apart. The injections consisted of 200 pmols of the liposomes complexed with 12.5 μ g of total DNA. The results are

shown in Table 4 below.

TABLE 4

Putative Anti-tumor Genes	Tumor Size(mm ³)
BAP	194.7 ± 11.9
TSPf	135.9 ± 11.9*
Laminin peptide	126.4 ± 8.4*
Angiostatin	95.2 ± 6.3*,**

* TSPf, Laminin peptide, and Angiostatin vs.

BAP, p<0.05

** Angiostatin vs. BAP, p<0.01

As shown in Table 4 above, the cationic liposomes containing only DNA encoding various anti-angiogenic peptides (TSPf, laminin peptide and angiostatin) significantly inhibited tumor growth.

Next, MCF7 cells (American Type Tissue Culture, Bethesda, MD), which are a breast cancer cell line with two normal p53 alleles, were evaluated as described above except that 4.0×10^6 cells were injected into the mice; and the third injection contained 12 µg of the DNA. Each injection was 10 days apart. Nine mice were injected with each of the following treatments except for regimen (1), in which 8 mice were treated: (1) untreated; (2) BAP; (3) p53; and (4) p53 + TSPf. The sizes of the tumors were measured 7 days after the third injection. The results are shown in Table 5 below.

TABLE 5

Effect of p53 and TSPf on MCF7s Cells	
Putative Anti-tumor Genes	Tumor Size (mm ³)
Untreated	124.6 ± 7.3
BAP	136 ± 16.8
p53	83.1 ± 13.6*
p53 + TSPf	69.0 ± 13**

* p53 vs. untreated or BAP, p<0.05

** p53 + TSPf vs. untreated or BAP, p<0.01

As shown in Table 5 above, the most effective therapy against MCF7s was p53 and TSPf. The significance level for the p53 + TSPf therapy was greater than for p53 alone when they were compared against either the untreated or the BAP groups.

The above experiment verifies that p53 and TSPf decreased the MCF7s tumor more than the p53 treated or the untreated groups. 4×10^5 MCF7 cells were injected bilaterally into the mammary fat pads of the 28 nude mice. After two weeks of growth, these mice were randomly divided into four groups: 1) empty vector, 2) p53, 3) p53 + TSPf, and 4) untreated. The mice received one injection of 200 pmoles of liposomes complexed with 14 µg of DNA, and the tumors from the various treatment groups were measured 10 days after the treatment. The results are shown in Table 6 below.

Table 6

Putative Anti-tumor Genes	Tumor Size (mm ³)
Empty vector-	54.7±4.0
p53	45.5±5.0
p53 + TSPf	33.9±3.6*
Untreated	61.9±8.3

*, p53 + TSPf vs Untreated, p<.025

As shown in Table 6, the additional reduction of the tumor by the combined use of p53 and TSPf (also in Tables 1, 2, and 4 above) compared to the use of p53 only, suggest that TSPf and p53 have different mechanisms of action. Although this does not preclude that the target of p53 is the vasculature of the tumor, the mechanism of inhibition of the tumor by p53 is not known at present. However, any mechanism of tumor inhibition by p53 and/or thrombospondin I must account for the low transfection efficiency of the tumor. Again, with a liposome complexed to a chloramphenicol acetyltransferase marker, it has been demonstrated that less than 5% of the tumor derived from MDA-MB-435 cells was transfected with the marker gene.

Besides p53 and the antiangiogenic fragment of thrombospondin I, we determined that liposomes complexed to DNA encoding the laminin peptide inhibits tumor growth. More specifically, after administering the anesthetic, Metofane, to 24 female athymic nude mice, the mice were injected with 3.0×10^5 MDA-MB-435 tumor cells into the mammary fat pad using a stepper and a 27.5 g needle. Two weeks later, the mice whose tumors grew were divided into various treatment regimens, 8 mice per each regimen. The treatment regimens were as follows: (1) BAP, (2) laminin, and (3) p53 + laminin. The mice were injected intravenously with 200 pmols of the liposomes complexed with 12.5 mgs of total DNA 6.25 mg of each vector when a combination was used. The mice then received 3 injections, each 10 days apart. The tumors were measured at the time of each injection and at the time of the last injection. The results are shown in Table 7 below.

TABLE 7

Putative Anti-tumor genes	Tumor Size (mm ³)
BAP	345 ± 23.5
Laminin peptide	280 ± 32.4
Laminin peptide + p53	192 ± 10.5*

* BAP vs. Laminin peptide + p53, p<0.05

As shown in Table 7 above, cationic liposomes containing a combination of DNAs encoding laminin peptide + p53 was unexpectedly more effective in reducing tumor growth than when DNA encoding the anti-angiogenic peptide was used alone. Thus, the addition of a tumor suppressor gene, p53, enhances the anti-tumor effect of the anti-angiogenic peptide.

Although intravenous injection are preferred, the method of administration of the liposome:DNA complex is not critical. In figure 1, it was found that intratumoral injections are effective, and it also supports that the therapy is effective against tumors other than breast cancer. In this experiment, 18 mice were injected with 3×10^5 C6 glioma cells (rat brain tumors) subcutaneously. Six days after the injections of these cells, the mice were separated into 3 groups: 1) BAP, 2) FLK-DN (a dominant negative receptor), and 3) angiostatin. After the second intratumoral injection, there was a statistical difference between the angiostatin and the BAP groups. Thus, this therapy is effective when given intratumorally and is effective as expected tumors other than breast tumors.

It was also found that this liposome: secretory angiostatin construct was more effective than the non-secreted analog. In brief, we injected 24 nude mice with 3×10^5 MDA-MB-435 cells inserted at the 5 prime end of the their construct. Two weeks later the mice were divided into three groups, they received the following therapies intravenously: 1) liposome:BAP, 2) liposome:secreted angiostatin, and 3) liposome:angiostatin. The concentration of DNA injected into the

mice was 14.5 ugs. The mice received one injection of the liposome:DNA complex and their tumors were measured 10 days after the injection.

Table 8

Efficacy of Secretory Angiostatin	
Therapeutic Genes	Tumor Size (mm ³)
Angiostatin	28.8±2.2
Angiostatin-Secretory	18.6±1.8*
BAP	30.5±3.3

*, $p < 0.05$, BAP vs. Angiostatin-secretory

As seen in table 8, the secretory angiostatin treatment group was much more effective than the empty vector control or the angiostatin treatment group in reducing the size of the tumor. From this experiment, it is evident that a secretory sequence inserted into the 5' portion of the antiangiogenic inhibitor will increase its efficacy.

In vitro assays indicate that cationic polymers will significantly improve the present therapy. When a carrier such as a cationic lipid was used in this *in vitro* assay, the inhibitory effect (of the genes p53, TSPf, and the combination of p53 and TSPf) was marginal whereas another vector, Superfect (a cationic polymer), was much more effective as a carrier. This is because Superfect was 15 times more effective than the cationic liposomes in transfecting endothelial cells with the CAT marker. The cationic liposomes used in this section was DOSPER (Boehringer), which of the 14 lipids tested gave the best results. Included in this panel of 14 lipids that we tested was lipofectin (BRL) which is a mixture of DOTMA/DOPE that we have used in an *in vivo* study. In brief, we plated 1×10^6 Huvec cells into each well of a 6 well plate. 25uls of Superfect complexed with 2 ugs of DNA was added to each plate 24 hours after the initial seeding of the cells. 36 hours after the transfection, the cells were lysed and the amount of CAT protein was assayed.

Table 9

Vectors	Activity(DPMs/protein)
Cationic liposomes with BAP	31.1±7.2
Cationic liposomes with CAT	682±129
Superfect with BAP	21.4±0.458
Superfect with CAT	10816±687*
$p < 0.001$, Superfect-CAT vs Cationic liposome-CAT	

This experiment clearly demonstrates that this cationic polymer is a superior in the transfection of endothelial cells, which is a likely target of the therapeutic gene. We have found that Superfect is a better transfection agent than cationic liposomes for many different cell lines. Since Superfect which is a cationic polymers is such an efficient carrier of DNA, this underscores possibility that non-viral carriers as a class of carriers will be effective in decreasing the tumor size. As a result, other non-viral carriers besides liposomes should be included in this patent.

Transfection of Huvec Cells with various inhibitors was as follows. 1.0×10^5 Huvec cells (Clonetics), a human endothelial cell line, were plated into each well of a 6-well plate, and placed in a CO₂ incubator at 37°C. Twenty-four hours later, the cells were transfected with 25 ml of Superfect (Qiagen), a cationic polymer, complexed to 2.0 mg of various DNA vectors, i.e., (1) BAP vector; (2) p53 vector; (3) TSPf vector; and (4) p53 vector + TSPf vector. After the cells were exposed for 2 hours to this complex at 37°C, the media was removed, and replaced with fresh EGM media (Clonetics, Inc.) containing 10% (v/v) fetal calf serum, and 1.0% (w/v) glutamine, and the cells placed in a CO₂ incubator at 37°C. Twenty-four hours later, the cell number in each plate was determined by the 3-(4,5-dimethylthiazol-2-yl)-(3-carboxymethoxyphenyl)-2,5-di-phenyltetra-zolium bromide (MTS) assay described by Butke et al, *J. Immunol. Methods*, 157:233-240 (1993).

The results are shown in Figure 1 attached hereto dealing with the intratumoral injections of liposome:DNA complexes and its effect on the tumor dimension 6 to 12 days after injection.

As shown in Figure 1, it was found that p53, TSPf, and the combination therapy of p53 and TSPf were effective at inhibiting endothelial cells *in vitro*. The combination of p53 and TSPf was the most effective at inhibiting endothelial

cells. There was a close correlation between the therapeutic genes reducing the tumor size *in vivo* and their effects on endothelial cell number *in vitro*.

Figure 2 shows the effect of different treatment groups on endothelial cells in vitro. it was found that the percentage of BAP control is significantly decreased when using BAP-p53 and BAPf-TSPf as tretment groups. A further synergistic decrease is achieved when using BAP-p53/TSPf a the treatment group.

While the invention has been described in detail, and with reference to specific embodiments thereof, it will be apparent to one of ordinary skill in the art that various changes and modifications can be made therein without departing from the spirit and scope thereof.

10

15

20

25

30

35

40

45

50

55

SEQUENCE LISTING

5 (1) GENERAL INFORMATION:

(i) APPLICANT:

(A) NAME: Archibald James MIXSON
 (B) STREET: 1 Baederwood Ct.
 10 (C) CITY: Rockville
 (D) STATE: Maryland
 (E) COUNTRY: USA
 (F) POSTAL CODE (ZIP): 21201

15 (ii) TITLE OF INVENTION: CATIONIC VEHICLE:DNA COMPLEXES AND THEIR
 USE IN GENE THERAPY

(iii) NUMBER OF SEQUENCES: 33

(iv) COMPUTER READABLE FORM:

20 (A) MEDIUM TYPE: Floppy disk
 (B) COMPUTER: IBM PC compatible
 (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 (D) SOFTWARE: PatentIn Release #1.0, Version #1.30 (EPO)

25 (v) CURRENT APPLICATION DATA:
 APPLICATION NUMBER: EP 97112154.6

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

30 (A) LENGTH: 217 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: linear

35 (ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

40 Met Thr Glu Asn Lys Glu Leu Ala Asn Glu Leu Arg Arg Pro Pro Leu
 1 5 10 15
 Cys Tyr His Asn Gly Val Gln Tyr Arg Asn Asn Glu Glu Trp Thr Val
 20 25 30
 45 Asp Ser Cys Thr Glu Cys His Cys Gln Asn Ser Val Thr Ile Cys Lys
 35 40 45
 Lys Val Ser Cys Pro Ile Met Pro Cys Ser Asn Ala Thr Val Pro Asp
 50 55 60
 Glu Glu Cys Cys Pro Arg Cys Trp Pro Ser Asp Ser Ala Asp Asp Gly
 65 70 75 80

55

Trp Ser Pro Trp Ser Glu Trp Thr Ser Cys Ser Thr Ser Cys Gly Asn
 85 90 95
 5 Gly Ile Gln Gln Arg Gly Arg Ser Cys Asp Ser Leu Asn Asn Arg Cys
 100 105 110
 Glu Gly Ser Ser Val Gln Thr Arg Thr Cys His Ile Gln Glu Cys Asp
 115 120 125
 10 Lys Arg Phe Lys Gln Asp Gly Gly Trp Ser His Trp Ser Pro Trp Ser
 130 135 140
 Ser Cys Val Thr Cys Gly Asp Gly Val Ile Thr Arg Ile Thr Leu Cys
 145 150 155 160
 15 Asn Ser Pro Ser Pro Gln Met Asn Gly Lys Pro Cys Glu Gly Glu Ala
 165 170 175
 Arg Glu Thr Lys Lys Ala Cys Lys Lys Asp Ala Cys Pro Ile Asn Glu
 180 185 190
 20 Gly Trp Gly Pro Trp Ser Pro Trp Asp Ile Cys Ser Val Thr Cys Gly
 195 200 205
 Gly Gly Val Gln Lys Arg Ser Arg Leu
 210 215
 25

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 656 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 30

(ii) MOLECULE TYPE: DNA (genomic)
 35

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

40 ATGACTGAAG AGAACAAGA GTTGGCCAAT GAGCTGAGGC GGCCTCCCCT ATGCTATCAC 60
 AACGGAGTTC AGTACAGAAA TAACGAGGAA TGGACTGTTG ATAGCTGCAC TGAGTGTACAC 120
 TGTCAAGAACT CAGTTACCAT CTGCAAAAAG GTGCTCTGCC CCATCATGCC CTGCTCCAAT 180
 45 GCCACAGTTC CTGATGGAGA ATGCTGTCCT CGCTGTTGGC CCAGCGACTC TGCGGACGAT 240
 GGCTGGTCTC CATGGTCCGA GTGGACCTCC TGTCTACGA GCTGTGGCAA TGAATTACAG 300
 50 CAGCGCGGCC GCTCCTGCGA TAGCCTCAAC AACCGATGTG AGGGCTCCTC GGTCCAGACA 360
 CGGACCTGCC ACATTACGGA GTGTGACAAA AGATTAAAC AGGATGGTGG CTGGAGCCAC 420

55

TGGTCCCCGT GGTCACTTGG TTCTGTGACA TGTGGTGATG GTGTGATCAC AAGGATCCGG 480
 CTCTGCAACT CTCCCAGCCC CCAGATGAAT GGGAAACCCT GTGAAGGCGA AGCGCGGGAG 540
 ACCAAAGCCT GCAAGAAAGA CGCTGCCCC ATCAATGGAG GCTGGGGTCC TTGGTCACCA 600
 TGGGACATCT GTTCTGTCAC CTGTGGAGGA GGGGTACAGA AACGTAGTCT CTCTAA 656

(2) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 439 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS:

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

Met	Thr	Glu	Asn	Lys	Glu	Leu	Ala	Asn	Glu	Leu	Arg	Arg	Pro	Pro	Leu
1				5					10					15	
Cys	Tyr	His	Asn	Gly	Val	Gln	Tyr	Arg	Asn	Asn	Glu	Glu	Trp	Thr	Val
			20					25					30		
Asp	Ser	Cys	Thr	Glu	Cys	His	Cys	Gln	Asn	Ser	Val	Thr	Ile	Cys	Lys
		35					40					45			
Lys	Val	Ser	Cys	Pro	Ile	Met	Pro	Cys	Ser	Asn	Ala	Thr	Val	Pro	Asp
	50					55					60				
Glu	Glu	Cys	Cys	Pro	Arg	Cys	Trp	Pro	Ser	Asp	Ser	Ala	Asp	Asp	Gly
65					70					75					80
Trp	Ser	Pro	Trp	Ser	Glu	Trp	Thr	Ser	Cys	Ser	Thr	Ser	Cys	Gly	Asn
			85					90					95		
Gly	Ile	Gln	Gln	Arg	Gly	Arg	Ser	Cys	Asp	Ser	Leu	Asn	Asn	Arg	Cys
			100					105					110		
Glu	Gly	Ser	Ser	Val	Gln	Thr	Arg	Thr	Cys	His	Ile	Gln	Glu	Cys	Asp
		115					120					125			
Lys	Arg	Phe	Lys	Gln	Asp	Gly	Gly	Trp	Ser	His	Trp	Ser	Pro	Trp	Ser
	130					135					140				
Ser	Cys	Val	Thr	Cys	Gly	Asp	Gly	Val	Ile	Thr	Arg	Ile	Thr	Leu	Cys
145					150					155					160
Asn	Ser	Pro	Ser	Pro	Gln	Met	Asn	Gly	Lys	Pro	Cys	Glu	Gly	Glu	Ala
				165					170				175		
Arg	Glu	Thr	Lys	Lys	Ala	Cys	Lys	Lys	Asp	Ala	Cys	Pro	Ile	Asn	Glu
			180					185					190		

Gly Trp Gly Pro Trp Ser Pro Trp Asp Ile Cys Ser Val Thr Cys Gly
 195 200 205
 5 Gly Gly Val Gln Lys Arg Ser Arg Leu Cys Val His Ser Arg Met Thr
 210 215 220
 Glu Asn Lys Glu Leu Ala Asn Glu Leu Arg Arg Pro Pro Leu Cys Tyr
 225 230 235 240
 10 His Asn Gly Val Gln Tyr Arg Asn Asn Glu Glu Trp Thr Val Asp Ser
 245 250 255
 Cys Thr Glu Cys His Cys Gln Asn Ser Val Thr Ile Cys Lys Lys Val
 260 265 270
 15 Ser Cys Pro Ile Met Pro Cys Ser Asn Ala Thr Val Pro Asp Glu Glu
 275 280 285
 Cys Cys Pro Arg Cys Trp Pro Ser Asp Ser Ala Asp Asp Gly Trp Ser
 290 295 300
 20 Pro Trp Ser Glu Trp Thr Ser Cys Ser Thr Ser Cys Gly Asn Gly Ile
 305 310 315 320
 Gln Gln Arg Gly Arg Ser Cys Asp Ser Leu Asn Asn Arg Cys Glu Gly
 325 330 335
 25 Ser Ser Val Gln Thr Arg Thr Cys His Ile Gln Glu Cys Asp Lys Arg
 340 345 350
 30 Phe Lys Gln Asp Gly Gly Trp Ser His Trp Ser Pro Trp Ser Ser Cys
 355 360 365
 Val Thr Cys Gly Asp Gly Val Ile Thr Arg Ile Thr Leu Cys Asn Ser
 370 375 380
 35 Pro Ser Pro Gln Met Asn Gly Lys Pro Cys Glu Gly Glu Ala Arg Glu
 385 390 395 400
 Thr Lys Lys Ala Cys Lys Lys Asp Ala Cys Pro Ile Asn Glu Gly Trp
 405 410 415
 40 Gly Pro Trp Ser Pro Trp Asp Ile Cys Ser Val Thr Cys Gly Gly Gly
 420 425 430
 Val Gln Lys Arg Ser Arg Leu
 435
 45

(2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:

- 50 (A) LENGTH: 1326 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

55

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

10	ATGACTGAAG AGAACAAAGA GTTGCCCAAT GAGCTGAGGC GGCCTCCCCCT ATGCTATCAC	60
	AACGGAGTTC AGTACAGAAA TAACGAGGAA TGGACTGTTG ATAGCTGCAC TGAGTGTCAC	120
	TGTCAGAACT CAGTTACCAT CTGCAAAAAG GTGTCCTGCC CCATCATGCC CTGCTCCAAT	180
15	GCCACAGTTC CTGATGGAGA ATGCTGTCCT CGCTGTTGGC CCAGCGACTC TGCGGACGAT	240
	GGCTGGTCTC CATGGTCCGA GTGGACCTCC TGTTCCTACGA GCTGTGGCAA TGGAAATTCAG	300
	CAGCGCGGCC GCTCCTGCGA TAGCCTCAAC AACCGATGTG AGGGCTCCTC GGTCCAGACA	360
20	CGGACCTGCC ACATTCAGGA GTGTGACAAA AGATTAAAC AGGATGGTGG CTGGAGCCAC	420
	TGGTCCCCGT GGTCACTTTG TTCTGTGACA TGTGGTGATG GTGTGATCAC AAGGATCCGG	480
25	CTCTGCAACT CTCCCAGCCC CCAGATGAAT GGGAAACCCT GTGAAGGCGA AGCGCGGGAG	540
	ACCAAAGCCT GCAAGAAAGA CGCCTGCCCC ATCAATGGAG GCTGGGGTCC TTGGTCACCA	600
	TGGGACATCT GTTCTGTGAC CTGTGGAGGA GGGGTACAGA AACGTAGTCG TCTCTGCGTC	660
30	GACTCTAGAA TGACTGAAGA GAACAAAGAG TTGGCCAATG AGCTGAGGCG GCCTCCCCCTA	720
	TGCTATCACA ACGGAGTTCA GTACAGAAAT AACGAGGAAT GGACTGTGA TAGCTGCACT	780
35	GAGTGTCACT GTCAGAACTC AGTTACCATC TGCAAAAAGG TGTCTGCCC CATCATGCCC	840
	TGCTCCAATG CCACAGTTCC TGATGGAGAA TGCTGTCTC GCTGTTGGCC CAGCGACTCT	900
	GCGGACGATG GCTGGTCTCC ATGGTCCGAG TGGACCTCCT GTTCTACGAG CTGTGGCAAT	960
40	GGAATTCAGC AGCGCGGCCG CTCCTGCGAT AGCCTCAACA ACCGATGTGA GGGCTCCTCG	1020
	GTCCAGACAC GGACCTGCCA CATTCAAGGAG TGTGACAAA GATTAAACA GGATGGTGGC	1080
	TGGAGCCACT GGTCCCCGTG GTCATCTTGT TCTGTGACAT GTGGTGATGG TGTGATCACA	1140
45	AGGATCCGGC TCTGCAACTC TCCCAGCCCC CAGATGAATG GGAAACCCTG TGAAGGCGAA	1200
	GCGCGGGAGA CCAAAGCCTG CAAGAAAGAC GCCTGCCCCA TCAATGGAGG CTGGGGTCCT	1260
50	TGGTCACCAT GGGACATCTG TTCTGTCACC TGTGGAGGAG GGGTACAGAA ACGTAGTCGT	1320
	CTCTAA	1326

55

(2) INFORMATION FOR SEQ ID NO: 5:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 6 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

Met Tyr Ile Gly Ser Arg
 1 5

(2) INFORMATION FOR SEQ ID NO: 6:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 33 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

GTCGACATGT ATATTGGTTC TCGTTAAGTC GAC 33

(2) INFORMATION FOR SEQ ID NO: 7:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 25 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

Met Tyr Ile Gly Ser Arg Gly Lys Ser Tyr Ile Gly Ser Arg Gly Lys
 1 5 10 15

Ser Tyr Ile Gly Ser Arg Gly Lys Ser
 20 25

(2) INFORMATION FOR SEQ ID NO: 8:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 90 base pairs

(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

5 (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

GTCGACATGT ATATTGGTTC TCGTGTA AAA GATATATTGG TTCTCGTGGT AAAAGATATA 60

TTGGTTCTCG TGGTAAAAGA TAAGTCGACC 90

15 (2) INFORMATION FOR SEQ ID NO: 9:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 13 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS:
(D) TOPOLOGY: linear

20 (ii) MOLECULE TYPE: protein

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

Met Leu Tyr Lys Lys Ile Ile Lys Lys Leu Leu Glu Ser
1 5 10

30 (2) INFORMATION FOR SEQ ID NO: 10:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 54 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

35 (ii) MOLECULE TYPE: DNA (genomic)

40 (iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

45 GTCGACATGC TTTATAAGAA GATCATCAAG AAGCTTCTTG AGAGTTAAGT CGAC 54

(2) INFORMATION FOR SEQ ID NO: 11:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 46 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS:
(D) TOPOLOGY: linear

50

55

(ii) MOLECULE TYPE: protein

5 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

Met Leu Tyr Lys Lys Ile Ile Lys Lys Leu Leu Glu Ser Gly Lys Ser
1 5 10 15

10 Leu Tyr Lys Lys Ile Ile Lys Lys Leu Leu Glu Ser Gly Lys Ser Leu
20 25 30

Tyr Lys Lys Ile Ile Lys Lys Leu Leu Glu Ser Gly Lys Ser
35 40 45

15 (2) INFORMATION FOR SEQ ID NO: 12:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 153 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
20 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

25 (iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:

GTCGACATGC TTTATAAGAA GATCATCAAG AAGCTTCTTG AGAGTGGTAA AAGACTTTAT 60
30 AAGAAGATCA TCAAGAAGCT TCTTGAGAGT GGTAAAAGAC TTTATAAGAA GATCATCAAG 120
AAGCTTCTTG AGAGTGGTAA AAGATAAGTC GAC 153

35 (2) INFORMATION FOR SEQ ID NO: 13:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 9 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS:
40 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

45 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:

Met Phe Cys Tyr Trp Lys Val Cys Trp
1 5

50 (2) INFORMATION FOR SEQ ID NO: 14:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 42 base pairs
(B) TYPE: nucleic acid

55

(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

5 (ii) MOLECULE TYPE: DNA (genomic)
(iii) HYPOTHETICAL: NO

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:

GTCGACATGT TCTTGTATTG GAAGGGATTG TGGTAAGTCG AC 42

(2) INFORMATION FOR SEQ ID NO: 15:

15 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 33 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS:
(D) TOPOLOGY: linear

20 (ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15:

25 Met Phe Cys Tyr Trp Lys Val Cys Trp Gly Lys Ser Phe Cys Tyr Trp
1 5 10 15

Lys Val Cys Trp Gly Lys Ser Phe Cys Trp Lys Val Cys Trp Gly Lys
20 25 30

30 Ser

(2) INFORMATION FOR SEQ ID NO: 16:

35 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 117 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

40 (ii) MOLECULE TYPE: DNA (genomic)
(iii) HYPOTHETICAL: NO

45 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 16:

GTCGACATGT TCTTGTATTG GAAGGGATTG TGGGGTAAAA GATTCTTGTA TTGGAAGGGA 60

TTGTGGGGTA AAAGATTCTT GTATTGGAAG GGATTGTGGG GTAAAAGATA AGTCGAC 117

50 (2) INFORMATION FOR SEQ ID NO: 17:

(i) SEQUENCE CHARACTERISTICS:

55

5 (A) LENGTH: 5 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS:
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 17:

Met Gly Arg Gly Asp
1 5

15 (2) INFORMATION FOR SEQ ID NO: 18:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 42 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
20 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 18:

GTCGACATGT CTTTGTCTTG GAAGACTTTG ACTTAAGTCG AC 42

30 (2) INFORMATION FOR SEQ ID NO: 19:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 22 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS:
35 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

40 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 19:

Met Gly Arg Gly Asp Gly Lys Ser Gly Arg Gly Asp Gly Lys Ser Gly
1 5 10 15

Arg Gly Asp Gly Lys Ser
20

45 (2) INFORMATION FOR SEQ ID NO: 20:

(i) SEQUENCE CHARACTERISTICS:
50 (A) LENGTH: 81 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

55

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 20:

GTCGACATGG GTCGTGGTGA TGGTAAAAGA GGTCGTGGTG ATGGTAAAAG AGGTCGTGGT 60
GATGGTAAAA GATAAGTCGA C 81

(2) INFORMATION FOR SEQ ID NO: 21:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 310 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 21:

Met	Val	Tyr	Leu	Ser	Glu	Cys	Lys	Thr	Gly	Ile	Gly	Asn	Gly	Tyr	Arg	1	5	10	15
Gly	Thr	Met	Ser	Arg	Thr	Lys	Ser	Gly	Val	Ala	Cys	Gln	Lys	Trp	Gly	20	25	30	
Ala	Thr	Phe	Pro	His	Val	Pro	Asn	Tyr	Ser	Pro	Ser	Thr	His	Pro	Asn	35	40	45	
Glu	Gly	Leu	Glu	Glu	Asn	Tyr	Cys	Arg	Asn	Pro	Asp	Asn	Asp	Glu	Gln	50	55	60	
Gly	Pro	Trp	Cys	Tyr	Thr	Thr	Asp	Pro	Asp	Lys	Arg	Tyr	Asp	Tyr	Cys	65	70	75	80
Asn	Ile	Pro	Glu	Cys	Glu	Glu	Glu	Cys	Met	Tyr	Cys	Ser	Gly	Glu	Lys	85	90	95	
Tyr	Glu	Gly	Lys	Ile	Ser	Lys	Thr	Met	Ser	Gly	Lys	Asp	Cys	Gln	Ala	100	105	110	
Trp	Asp	Ser	Gln	Ser	Pro	His	Ala	His	Gly	Tyr	Ile	Pro	Ala	Lys	Phe	115	120	125	
Pro	Ser	Lys	Asn	Leu	Lys	Met	Asn	Tyr	Cys	His	Asn	Pro	Asp	Gly	Glu	130	135	140	
Pro	Arg	Pro	Trp	Cys	Phe	Thr	Thr	Asp	Pro	Thr	Lys	Arg	Trp	Glu	Tyr	145	150	155	160

Cys Asp Ile Pro Arg Cys Thr Thr Pro Pro Pro Pro Ser Pro Thr
 165 170 175
 5 Tyr Gln Cys Leu Lys Gly Arg Gly Glu Asn Tyr Arg Gly Thr Val Ser
 180 185 190
 Val Thr Val Ser Gly Lys Thr Cys Gln Arg Trp Ser Glu Gln Thr Pro
 195 200 205
 10 His Arg His Asn Arg Thr Pro Glu Asn Phe Pro Cys Arg Asn Leu Glu
 210 215 220
 Glu Asn Tyr Cys Arg Asn Pro Asp Gly Glu Thr Ala Pro Trp Cys Tyr
 225 230 235 240
 15 Thr Thr Asp Ser Gln Leu Arg Trp Glu Tyr Cys Glu Ile Pro Ser Cys
 245 250 255
 Glu Ser Ser Ala Ser Pro Asp Gln Ser Asp Ser Ser Val Pro Pro Glu
 260 265 270
 20 Glu Gln Thr Pro Val Val Gln Glu Cys Tyr Gln Ser Asp Gly Gln Ser
 275 280 285
 Tyr Arg Gly Thr Ser Ser Thr Thr Ile Thr Gly Lys Lys Cys Gln Ser
 290 295 300
 25 Glu Gln Thr Pro His Arg
 305 310

(2) INFORMATION FOR SEQ ID NO: 22:

30

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 645 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 35 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

40

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 22:

GTCGACATGG TGTATCTGTC AGAATGTAAG ACCGGCATCG GCAACGGCTA CAGAGGAACC 60
 45 ATGTCCAGGA CAAAGAGTGG TGTTCCTGT CAAAAGTGGG GTGCCACGTT CCCCCACGTA 120
 CCCAACTACT CTCCCAGTAC ACATCCCAAT GAGGGACTAG AAGAGAACTA CTGTAGGAAC 180
 CCAGACAATG ATGAACAAGG GCCTTGGTGC TACACTACAG ATCCGGACAA GAGATATGAC 240
 50 TACTGCAACA TTCCTGAATG TGAAGAGGAA TGCATGTACT GCAGTGGAGA AAAGTATGAG 300
 GGCAAAATCT CCAAGACCAT GTCTGGACTT GACTGCCAGG CCTGGGATTC TCAGAGCCCA 360

55

CATGCTCATG GATACATCCC TGCCAAATTT CCAAGCAAGA ACCTGAAGAT GAATTATTGC 420
 CACAACCCCTG ACGGGGAGCC AAGGCCCTGG TGCTTCACAA CAGACCCAC CAAACGCTGG 480
 GAATACTGTG ACATCCCCCG CTGCACAACA CCCCCGCCCC CACCCAGCCC AACCTACCAA 540
 TGTCTGAAAG GAAGAGGTGA AAATTACCGA GGGACCGTGT CTGTCACCGT GTCTGGGAAA 600
 ACCTGTCAGC GCTGGAGTGA GCAAACCCCT CATAGGTGAG TCGAC 645

(2) INFORMATION FOR SEQ ID NO: 23:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 623 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 23:

Met	Val	Tyr	Leu	Ser	Glu	Cys	Lys	Thr	Gly	Ile	Gly	Asn	Gly	Tyr	Arg
1				5					10					15	
Gly	Thr	Met	Ser	Arg	Thr	Lys	Ser	Gly	Val	Ala	Cys	Gln	Lys	Trp	Gly
			20					25					30		
Ala	Thr	Phe	Pro	His	Val	Pro	Asn	Tyr	Ser	Pro	Ser	Thr	His	Pro	Asn
			35				40					45			
Glu	Gly	Leu	Glu	Glu	Asn	Tyr	Cys	Arg	Asn	Pro	Asp	Asn	Asp	Glu	Gln
		50				55					60				
Gly	Pro	Trp	Cys	Tyr	Thr	Thr	Asp	Pro	Asp	Lys	Arg	Tyr	Asp	Tyr	Cys
	65				70					75				80	
Asn	Ile	Pro	Glu	Cys	Glu	Glu	Glu	Cys	Met	Tyr	Cys	Ser	Gly	Glu	Lys
				85					90					95	
Tyr	Glu	Gly	Lys	Ile	Ser	Lys	Thr	Met	Ser	Gly	Lys	Asp	Cys	Gln	Ala
			100					105					110		
Trp	Asp	Ser	Gln	Ser	Pro	His	Ala	His	Gly	Tyr	Ile	Pro	Ala	Lys	Phe
		115					120						125		
Pro	Ser	Lys	Asn	Leu	Lys	Met	Asn	Tyr	Cys	His	Asn	Pro	Asp	Gly	Glu
		130				135						140			
Pro	Arg	Pro	Trp	Cys	Phe	Thr	Thr	Asp	Pro	Thr	Lys	Arg	Trp	Glu	Tyr
	145				150					155				160	
Cys	Asp	Ile	Pro	Arg	Cys	Thr	Thr	Pro	Pro	Pro	Pro	Ser	Pro	Thr	
				165				170					175		

Tyr Gln Cys Leu Lys Gly Arg Gly Glu Asn Tyr Arg Gly Thr Val Ser
 180 185 190
 5 Val Thr Val Ser Gly Lys Thr Cys Gln Arg Trp Ser Glu Gln Thr Pro
 195 200 205
 His Arg His Asn Arg Thr Pro Glu Asn Phe Pro Cys Arg Asn Leu Glu
 210 215 220
 10 Glu Asn Tyr Cys Arg Asn Pro Asp Gly Glu Thr Ala Pro Trp Cys Tyr
 225 230 235 240
 Thr Thr Asp Ser Gln Leu Arg Trp Glu Tyr Cys Glu Ile Pro Ser Cys
 245 250 255
 15 Glu Ser Ser Ala Ser Pro Asp Gln Ser Asp Ser Ser Val Pro Pro Glu
 260 265 270
 Glu Gln Thr Pro Val Val Gln Glu Cys Tyr Gln Ser Asp Gly Gln Ser
 275 280 285
 20 Tyr Arg Gly Thr Ser Ser Thr Thr Ile Thr Gly Lys Lys Cys Gln Ser
 290 295 300
 Glu Gln Thr Pro His Arg Gly Lys Ser Met Val Tyr Leu Ser Glu Cys
 305 310 315 320
 25 Lys Thr Gly Ile Gly Asn Gly Tyr Arg Gly Thr Met Ser Arg Thr Lys
 325 330 335
 Ser Gly Val Ala Cys Gln Lys Trp Gly Ala Thr Phe Pro His Val Pro
 340 345 350
 30 Asn Tyr Ser Pro Ser Thr His Pro Asn Glu Gly Leu Glu Glu Asn Tyr
 355 360 365
 35 Cys Arg Asn Pro Asp Asn Asp Glu Gln Gly Pro Trp Cys Tyr Thr Thr
 370 375 380
 Asp Pro Asp Lys Arg Tyr Asp Tyr Cys Asn Ile Pro Glu Cys Glu Glu
 385 390 395 400
 40 Glu Cys Met Tyr Cys Ser Gly Glu Lys Tyr Glu Gly Lys Ile Ser Lys
 405 410 415
 Thr Met Ser Gly Lys Asp Cys Gln Ala Trp Asp Ser Gln Ser Pro His
 420 425 430
 45 Ala His Gly Tyr Ile Pro Ala Lys Phe Pro Ser Lys Asn Leu Lys Met
 435 440 445
 Asn Tyr Cys His Asn Pro Asp Gly Glu Pro Arg Pro Trp Cys Phe Thr
 450 455 460
 50 Thr Asp Pro Thr Lys Arg Trp Glu Tyr Cys Asp Ile Pro Arg Cys Thr
 55

465 470 475 480
 Thr Pro Pro Pro Pro Pro Ser Pro Thr Tyr Gln Cys Leu Lys Gly Arg
 5 485 490 495
 Gly Glu Asn Tyr Arg Gly Thr Val Ser Val Thr Val Ser Gly Lys Thr
 500 505 510
 Cys Gln Arg Trp Ser Glu Gln Thr Pro His Arg His Asn Arg Thr Pro
 10 515 520 525
 Glu Asn Phe Pro Cys Arg Asn Leu Glu Glu Asn Tyr Cys Arg Asn Pro
 530 535 540
 Asp Gly Glu Thr Ala Pro Trp Cys Tyr Thr Thr Asp Ser Gln Leu Arg
 15 545 550 555 560
 Trp Glu Tyr Cys Glu Ile Pro Ser Cys Glu Ser Ser Ala Ser Pro Asp
 565 570 575
 Gln Ser Asp Ser Ser Val Pro Pro Glu Glu Gln Thr Pro Val Val Gln
 20 580 585 590
 Glu Cys Tyr Gln Ser Asp Gly Gln Ser Tyr Arg Gly Thr Ser Ser Thr
 595 600 605
 Thr Ile Thr Gly Lys Lys Cys Gln Ser Glu Gln Thr Pro His Arg
 25 610 615 620

(2) INFORMATION FOR SEQ ID NO: 24:

30

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1284 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

35

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

40

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 24:

GTCGACATGG TGTATCTGTC AGAATGTAAG ACCGGCATCG GCAACGGCTA CAGAGGAACC 60
 ATGTCCAGGA CAAAGAGTGG TGTGCTGT CAAAAGTGGG GTGCCACGTT CCCCCACGTA 120
 45 CCAACTACT CTCCCAGTAC ACATCCCAAT GAGGGACTAG AAGAGAACTA CTGTAGGAAC 180
 CCAGACAATG ATGAACAAGG GCCTTGGTGC TACTACTACAG ATCCGGACAA GAGATATGAC 240
 TACTGCAACA TTCCTGAATG TGAAGAGGAA TGCATGTACT GCAGTGGAGA AAAGTATGAG 300
 50 GGCAAAATCT CCAAGACCAT GTCTGGACTT GACTGCCAGG CCTGGGATTC TCAGAGCCCA 360

55

5 CATGCTCATG GATACATCCC TGCCAAATTT CCAAGCAAGA ACCTGAAGAT GAATTATTGC 420
 CACAACCCCTG ACGGGGAGCC AAGGCCCTGG TGCTTCACAA CAGACCCAC CAAACGCTGG 480
 GAATACTGTG ACATCCCCCG CTGCACAACA CCCCCGCCCC CACCCAGCCC AACCTACCAA 540
 TGTCTGAAAG GAAGAGGTGA AAATTACCGA GGGACCGTGT CTGTCACCGT GTCTGGGAAA 600
 10 ACCTGTCAGC GCTGGAGTGA GCAAACCCCT CATAGGGGTA AAAGAATGGT GTATCTGTCA 660
 GAATGTAAGA CCGGCATCGG CAACGGCTAC AGAGGAACCA TGTCCAGGAC AAAGAGTGGT 720
 GTTGCCCTGTC AAAAGTGGGG TGCCACGTTT CCCCACGTAC CCAACTACTC TCCCAGTACA 780
 15 CATCCCAATG AGGGACTAGA AGAGAACTAC TGAGGAACC CAGACAATGA TGAACAAGGG 840
 CCTTGGTGCT AACTACAGA TCCGGACAAG AGATATGACT ACTGCAACAT TCCTGAATGT 900
 GAAGAGGAAT GCATGTACTG CAGTGGAGAA AAGTATGAGG GCAAAATCTC CAAGACCATG 960
 20 TCTGGACTTG ACTGCCAGGC CTGGGATTCT CAGAGCCCAC ATGCTCATGG ATACATCCCT 1020
 GCCAAATTTT CAAGCAAGAA CCTGAAGATG AATTATTGCC ACAACCTGA CGGGGAGCCA 1080
 25 AGGCCCTGGT GCTTCACAAC AGACCCACC AAACGCTGGG AATACTGTGA CATCCCCGC 1140
 TGCACAACAC CCCCCCCCC ACCCAGCCCA ACCTACCAAT GTCTGAAAGG AAGAGGTGAA 1200
 AATTACCGAG GGACCGTGTG TGTCACCGTG TCTGGGAAAA CCTGTCAGCG CTGGAGTGAG 1260
 30 CAAACCCCTC ATAGGTGAGT CGAC 1284

(2) INFORMATION FOR SEQ ID NO: 25:

35 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 125 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: linear

40 (ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 25:

45 Met Leu Pro Ile Cys Pro Gly Gly Ala Ala Arg Cys Gln Val Thr Leu
 1 5 10 15
 Arg Glu Leu Phe Asp Arg Ala Val Val Leu Ser His Tyr Ile His Asn
 20 25 30
 50 Leu Ser Ser Glu Met Phe Ser Glu Phe Glu Lys Arg Tyr Thr His Gly
 35 40 45
 Arg Gly Phe Ile Thr Lys Ala Ile Asn Ser Cys His Thr Ser Ser Leu
 50 55 60

55

Ala Thr Pro Glu Asp Lys Glu Gln Ala Gln Gln Met Asn Gln Lys Asp
65 70 75 80

5 Phe Leu Ser Leu Ile Val Ser Ile Leu Arg Ser Trp Asn Glu Pro Leu
85 90 95

Tyr His Leu Val Thr Glu Val Arg Gly Met Gln Glu Ala Pro Gln Ala
100 105 110

10 Ile Leu Ser Lys Ala Val Glu Ile Glu Glu Gln Thr Lys
115 120 125

(2) INFORMATION FOR SEQ ID NO: 26:

15 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 387 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
20 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 26:

GTGACATGT TGCCCATCTG TCCCGGCGGG GCTGCCCCGAT GCCAGGTGAC CCTTCGAGAC 60
30 CTGTTTGACC GCGCCGTCGT CCTGTCCCAC TACATCCATA ACCTCTCCTC AGAAATGTTT 120
AGCGAATTCG ATAAACGGTA TACCCATGGC CGGGGGTTCA TTACCAAGGC CATCAACAGC 180
TGCCACACTT CTTCCTTGC CACCCCCGAA GACAAGGAGC AAGCCCAACA GATGAATCAA 240
35 AAAGACTTTC TGAGCCTGAT AGTCAGCATA TTGCGATCCT GGAATGAGCC TCTGTATCAT 300
CTGGTCACGG AAGTACGTGG TATGCAAGAA GCCCGGAGG CTATCCTATC CAAAGCTGTA 360
40 GAGATTGAGG AGCAAACCTA AGTCGAC 387

(2) INFORMATION FOR SEQ ID NO: 27:

45 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 255 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS:
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

50

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 27:

55

Met Leu Pro Ile Cys Pro Gly Gly Ala Ala Arg Cys Gln Val Thr Leu
 1 5 10 15
 5 Arg Glu Leu Phe Asp Arg Ala Val Val Leu Ser His Tyr Ile His Asn
 20 25 30
 Leu Ser Ser Glu Met Phe Ser Glu Phe Glu Lys Arg Tyr Thr His Gly
 35 40 45
 10 Arg Gly Phe Ile Thr Lys Ala Ile Asn Ser Cys His Thr Ser Ser Leu
 50 55 60
 Ala Thr Pro Glu Asp Lys Glu Gln Ala Gln Gln Met Asn Gln Lys Asp
 65 70 75 80
 15 Phe Leu Ser Leu Ile Val Ser Ile Leu Arg Ser Trp Asn Glu Pro Leu
 85 90 95
 Tyr His Leu Val Thr Glu Val Arg Gly Met Gln Glu Ala Pro Gln Ala
 100 105 110
 20 Ile Leu Ser Lys Ala Val Glu Ile Glu Glu Gln Thr Lys Gly Lys Ser
 115 120 125
 Met Leu Pro Ile Cys Pro Gly Gly Ala Ala Arg Cys Gln Val Thr Leu
 130 135 140
 25 Arg Glu Leu Phe Asp Arg Ala Val Val Leu Ser His Tyr Ile His Asn
 145 150 155 160
 Leu Ser Ser Glu Met Phe Ser Glu Phe Glu Lys Arg Tyr Thr His Gly
 165 170 175
 Arg Gly Phe Ile Thr Lys Ala Ile Asn Ser Cys His Thr Ser Ser Leu
 180 185 190
 35 Ala Thr Pro Glu Asp Lys Glu Gln Ala Gln Gln Met Asn Gln Lys Asp
 195 200 205
 Phe Leu Phe Leu Ser Leu Ile Val Ser Ile Leu Arg Ser Trp Asn Glu
 210 215 220
 40 Pro Leu Tyr His Leu Val Thr Glu Val Arg Gly Met Gln Glu Ala Pro
 225 230 235 240
 Gln Ala Ile Leu Ser Lys Ala Val Glu Ile Glu Glu Gln Thr Lys
 245 250 255
 45

(2) INFORMATION FOR SEQ ID NO: 28:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 756 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 28:

ATGTTGCCCA TCTGTCCCGG CGGGGCTGCC CGATGCCAGG TGACCCCTCG AGACCTGTTT 60
 10 GACCGCGCCG TCGTCCTGTC CCACTACATC CATAACCTCT CCTCAGAAAT GTTCAGCGAA 120
 TTCGATAAAC GGTATACCCA TGGCCGGGGG TTCATTACCA AGGCCATCAA CAGCTGCCAC 180
 15 ACTTCTTCCC TTGCCACCCC CGAAGACAAG GAGCAAGCCC AACAGATGAA TCAAAAAGAC 240
 TTTCTGAGCC TGATAGTCAG CATATTGCGA TCCTGGAATG AGCCTCTGTA TCATCTGGTC 300
 ACGGAAGTAC GTGGTATGCA AGAAGCCCCG GAGGCTATCC TATCCAAAGC TGTAGAGATT 360
 20 GAGGAGCAAA CCGGTAAAAG AATGTTGCCC ATCTGTCCCG GCGGGGCTGC CCGATGCCAG 420
 GTGACCCCTC GAGACCTGTT TGACCGCGCC GTCGTCTGT CCCACTACAT CCATAACCTC 480
 TCCTCAGAAA TGTTCAGCGA ATTCGATAAA CGGTATACCC ATGGCCGGGG GTTCATTACC 540
 25 AAGGCCATCA ACAGCTGCCA CACTTCTTCC CTGCCCACCC CCGAAGACAA GGAGCAAGCC 600
 CAACAGATGA ATCAAAAAGA CTTTCTGAGC CTGATAGTCA GCATATTGCG ATCCTGGAAT 660
 GAGCCTCTGT ATCATCTGGT CACGGAAGTA CGTGGTATGC AAGAAGCCCC GGAGGCTATC 720
 30 CTATCCAAAG CTGTAGAGAT TGAGGAGCAA ACCTAA 756

(2) INFORMATION FOR SEQ ID NO: 29:

35

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 161 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

40

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

45

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 29:

ATGCTGAGGC GGCCTCCCCT ATGCTATCAC AACGGAGTTC AGTACAGAAA TAACGGTAAA 60
 50 AGATCCCCGT GGTCACTCTG TTCTGTGACA TGTGGTGATG GTGTGATGGT AAAAGAAGTG 120
 GTACCCTGTA GACAAGACAG TGGACACCTC CTCCTCATTA A 161

55

(2) INFORMATION FOR SEQ ID NO: 30:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 63 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 30:

```

Met Leu Arg Arg Pro Pro Leu Cys Tyr His Asn Gly Val Gln Tyr Arg
1          5          10          15

Asn Asn Glu Glu Trp Thr Val Asp Ser Gly Lys Ser Ser Pro Trp Ser
          20          25          30

Ser Cys Ser Val Thr Cys Gly Asp Gly Val Ile Thr Arg Ile Gly Lys
          35          40          45

Ser Ser Pro Trp Asp Ile Cys Ser Val Thr Cys Gly Gly Gly Val
          50          55          60

```

(2) INFORMATION FOR SEQ ID NO: 31:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 185 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 31:

```

ATGCTGAGGC GGCCTCCCCT ATGCTATCAC AACGGAGTTC AGTACAGAAA TAACGGTAAA      60
AGATCCCCGT GGTCACTCTG TTCTGTGACA TGTGGTGATG GTGTGATGGT AAAAGAAGTG      120
GTACCCTGTA GACAAGACAG TGGACACCTC CTCCCCATTA TATTGTTTCT CGTGGTAAAA      180
GATAA                                           185

```

(2) INFORMATION FOR SEQ ID NO: 32:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 31 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 32:

TAGGTCTAGA ATGACTGAAG AGAACAAGA G

31

(2) INFORMATION FOR SEQ ID NO: 33:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 31 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 33:

ATGGTCTAGA TTAGAGACGA CTACGTTTCT G

31

Claims

1. A cationic liposome:DNA complex comprising DNA encoding an anti-angiogenic peptide.
2. A cationic polymer:DNA complex comprising the DNA encoding an antiangiogenic peptide.
3. The complex of Claim 1, wherein said complex additionally comprises DNA encoding a tumor suppressor protein.
4. The complex of Claim 2, wherein said complex additionally comprises DNA encoding a tumor suppressor protein.
5. The complex of Claim 1, wherein said cationic liposomes are comprised of one cationic lipid (i.e.- 1,2-dioleoyl-sn-glycero-3-ethylphosphocholine, 1,2-dimyristoyl-sn-glycero-3-ethylphosphocholine, and 2,3-diol-eyloxy)propyl-N,N,N- trimethyl-ammonium chloride) and may also be comprised of polyethylene glycol (i.e., a pegylated lipid-1,2-diacy-sn-glycero-3-phospho-ethanolamine-N-Poly(ethylene glycol) 2000) and a targeted ligand (the amino acid peptide RGD, ferritin, or antibodies targeted toward to HER2).
6. The complex of Claim 1, wherein said cationic liposomes are comprised of one cationic polymer polyethylenimine Polycat57, polylysine, polyhisitidine, and Superfect) and may also be comprised of polyethylene glycol and a targeted ligand (the amino acid peptide RGD, ferritin, or antibodies targeted toward to HER2, CEA).
7. The complex of Claims 4 or 5, wherein said tumor suppressor protein is selected from the group consisting of the p53, the p21 and the rb.
8. The complex of Claims 3 or 4, wherein said tumor suppressor protein is p53.
9. The complex of Claims 1, 2, or 3, wherein said anti-angiogenic peptide is selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID

NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, and SEQ ID NO:30.

10. The complex of Claims 1 or 3, wherein said DNA encoding an anti-angiogenic peptide is present in an amount of from 0.0025 to 0.16 $\mu\text{g}/\text{pM}$ of liposome.
11. The complex of Claims 2 or 4, wherein said DNA encoding an anti-angiogenic peptide is present in an amount of from 0.016 to 0.33 $\mu\text{g}/\mu\text{g}$ of polymer.
12. The complex of Claim 8, wherein said DNA encoding a tumor suppressor protein is present in an amount of from 0.0025 to 0.16 $\mu\text{g}/\text{pM}$.
13. The complex of Claim 8, wherein said DNA encoding a tumor suppressor protein is present in an amount of from 0.016 to 0.33 $\mu\text{g}/\text{pM}$.
14. Use of a cationic polymer:DNA complex comprising DNA encoding an anti-angiogenic peptide for the production of a medicament for inhibiting tumor growth in a subject which preferably comprises administering the same to a tumor-bearing subject.
15. The use of Claim 14, where said complex additionally comprises DNA encoding a tumor suppressor protein.
16. The use of claim 1, wherein said cationic liposome (i.e.-1,2-dioleoyl-sn-glycero-3-ethylphosphocholine, 1,2-dimyr-istoyl-sn-glycero-3-ethylphosphocholine, and 2,3-diol-eyloxy)propyl-N,N,N- trimethylammonium chloride and may also be comprised of polyethylene glycol (i.e., a pegylated lipid-1,2 -diacy-sn-glycero-3-phospho-ethanolamine-N-[Poly-(ethylene glycol) 2000) and a targeted ligand (the amino acid peptide RGD, ferritin, or antibodies targeted toward to HER2).
17. The use of claim 2, wherein said cationic polymer (i.e. (polyethytenimine Polycat57, polylysine, polyhisitidine, and Superfect) and may also be comprised of polyethylene glycol and a targeted ligand (the amino acid peptide RGD, ferritin, or antibodies targeted toward to HER2).
18. The use of Claim 15, wherein said tumor suppressor protein is selected from the group consisting of p53, the p21 and the rb.
19. The use of Claim 18, wherein said tumor suppressor protein is p53.
20. The use of Claim 14, wherein said anti-angiogenic peptide is selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, and SEQ ID NO:30.
21. The use of Claim 14 or Claim 15, wherein said DNA encoding an anti-angiogenic peptide is present in an amount from 0.0025 to 0.16 $\mu\text{g}/\text{pM}$ of liposome or 0.016 to 0.33 $\mu\text{g}/\mu\text{g}$ of polymer.
22. The use of Claims 14 or 15, wherein said DNA encoding a tumor suppressor protein is present in an amount of from 0.0025 to 0.16 $\mu\text{g}/\text{pM}$ of liposome or 0.016 to 0.33 $\mu\text{g}/\mu\text{g}$ of polymer.
23. A cationic polymer:DNA complex comprising a plasmid with one or more promoters expressing either the same gene product (SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, and SEQ ID NO:30) or combinations of these gene products.
24. A cationic liposome:DNA complex comprising a plasmid with an intervening internal ribosomal entry site sequence between two genes (SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:22, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, and SEQ ID NO:29) that are either the same or different.
25. The complex of Claim 1 or Claim 2, wherein said anti-angiogenic protein is secretory from the group consisting of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID

EP 0 819 758 A2

NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, and SEQ ID NO:30.

5

10

15

20

25

30

35


40

45

50


55

Figure 1
Intratumoral Injections of Liposome:DNA Complexes



*, Angio vs. BAP, $p < 0.05$

Figure 2
Effect of Different Treatment Groups on Endothelial
Cells *In Vitro*.



*- BAP vs BAP-p53 or BAP-TSPf, $p < 0.05$

** - BAP-p53 or BAP-TSPf vs BAP-p53/BAP-TSPf, $p < 0.01$